

International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

Climate Change: Human Impact on the Environment

Satya Dev

Lecturer in Geography, MSJ Govt. College, Bharatpur, Rajasthan, India

ABSTRACT: Global temperatures rose about 1.98°Foffsite link (1.1°C) from 1901 to 2008, but climate change refers to more than an increase in temperature. It also includes sea level rise, changes in weather patterns like drought and flooding, and much more. Things that we depend upon and value — water, energy, transportation, wildlife, agriculture, ecosystems, and human health — are experiencing the effects of a changing climate. The impacts of climate change on different sectors of society are interrelated. Drought can harm food production and human health. Flooding can lead to disease spread and damages to ecosystems and infrastructure. Human health issues can increase mortality, impact food availability, and limit worker productivity. Climate change impacts are seen throughout every aspect of the world we live in. However, climate change impacts are uneven across the country and the world — even within a single community, climate change impacts can differ between neighborhoods or individuals. Long-standing socioeconomic inequities can make underserved groups, who often have the highest exposure to hazards and the fewest resources to respond, more vulnerable.

KEYWORDS: climate change, human impact, environment, health, mortality, socioeconomic, hazards

I.INTRODUCTION

Humans impact the physical environment in many ways: overpopulation, pollution, burning fossil fuels, and deforestation. Changes like these have triggered climate change, soil erosion, poor air quality, and undrinkable water. These negative impacts can affect human behavior and can prompt mass migrations or battles over clean water.

Human impact on the environment (or anthropogenic impact) refers to changes to biophysical environments^[1] and to ecosystems, biodiversity, and natural resources^[2] caused directly or indirectly by humans. Modifying the environment to fit the needs of society is causing severe effects^{[3][4]} including global warming,^{[1][5]} environmental degradation^[1] (such as ocean acidification^{[1][6]}), mass extinction and biodiversity loss,^{[7][8][9][10]} ecological crisis, and ecological collapse. Some human activities that cause damage (either directly or indirectly) to the environment on a global scale include population growth,^{[11][12][13]} rapid economic growth,^[14] overconsumption, overexploitation, pollution, and deforestation. Some of the problems, including global warming and biodiversity loss, have been proposed as representing catastrophic risks to the survival of the human species.^{[15][16]}

The term anthropogenic designates an effect or object resulting from human activity. The term was first used in the technical sense by Russian geologist Alexey Pavlov, and it was first used in English by British ecologist Arthur Tansley in reference to human influences on climax plant communities. The atmospheric scientist Paul Crutzen introduced the term "Anthropocene" in the mid-1970s. The term is sometimes used in the context of pollution produced from human activity since the start of the Agricultural Revolution but also applies broadly to all major human impacts on the environment. [19][20][21] Many of the actions taken by humans that contribute to a heated environment stem from the burning of fossil fuel from a variety of sources, such as: electricity, cars, planes, space heating, manufacturing, or the destruction of forests. [22]

Overconsumption is a situation where resource use has outpaced the sustainable capacity of the ecosystem. It can be measured by the ecological footprint, a resource accounting approach which compares human demand on ecosystems



International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

with the amount of planet matter ecosystems can renew. Estimates indicate that humanity's current demand is 70% [24] higher than the regeneration rate of all of the planet's ecosystems combined. A prolonged pattern of overconsumption leads to environmental degradation and the eventual loss of resource bases.

Humanity's overall impact on the planet is affected by many factors, not just the raw number of people. Their lifestyle (including overall affluence and resource use) and the pollution they generate (including carbon footprint) are equally important. In 2008, The New York Times stated that the inhabitants of the developed nations of the world consume resources like oil and metals at a rate almost 32 times greater than those of the developing world, who make up the majority of the human population. [25]

Human civilization has caused the loss of 83% of all wild mammals and half of plants.^[26] The world's chickens are triple the weight of all the wild birds, while domesticated cattle and pigs outweigh all wild mammals by 14 to 1.^{[27][28]} Global meat consumption is projected to more than double by 2050, perhaps as much as 76%, as the global population rises to more than 9 billion, which will be a significant driver of further biodiversity loss and increased Greenhouse gas emissions.^{[29][30]}

Some scholars, environmentalists and advocates have linked human population growth or population size as a driver of environmental issues, including some suggesting this indicates an overpopulation scenario. [11] In 2011, over 15,000 scientists around the world issued a second warning to humanity which asserted that rapid human population growth is the "primary driver behind many ecological and even societal threats." [33] According to the Global Assessment Report on Biodiversity and Ecosystem Services, released by the United Nations' Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services in 2009, human population growth is a significant factor in contemporary biodiversity loss. [34] A 2007 report in Frontiers in Conservation Science proposed that population size and growth are significant factors in biodiversity loss, soil degradation and pollution. [35][36]

Some scientists and environmentalists, including Pentti Linkola, [37] Jared Diamond and E. O. Wilson, posit that human population growth is devastating to biodiversity. Wilson for example, has expressed concern when Homo sapiens reached a population of six billion their biomass exceeded that of any other large land dwelling animal species that had ever existed by over 100 times. [38]

However, attributing overpopulation as a cause of environmental issues is controversial. Demographic projections indicate that population growth is slowing and world population will peak in the 21st century, [31] and many experts believe that global resources can meet this increased demand, suggesting a global overpopulation scenario is unlikely. Other projections have the population continuing to grow into the next century. While some studies, including the British government's 2007 Economics of Biodiversity review, posit that population growth and overconsumption are interdependent, critics suggest blaming overpopulation for environmental issues can unduly blame poor populations in the Global South or oversimplify more complex drivers, leading some to treat overconsumption as a separate issue.

Advocates for further reducing fertility rates, among them Rodolfo Dirzo and Paul R. Ehrlich, argue that this reduction should primarily affect the "overconsuming wealthy and middle classes," with the ultimate goal being to shrink "the scale of the human enterprise" and reverse the "growthmania" which they say threatens biodiversity and the "life-support systems of humanity." [46]

The environmental impact of agriculture varies based on the wide variety of agricultural practices employed around the world. Ultimately, the environmental impact depends on the production practices of the system used by farmers. The connection between emissions into the environment and the farming system is indirect, as it also depends on other climate variables such as rainfall and temperature. There are two types of indicators of environmental impact: "means-based", which is based on the farmer's production methods, and "effect-based", which is the impact that farming methods have on the farming system or on emissions to the environment. An example of a means-based indicator would be the quality of groundwater that is affected by the amount of nitrogen applied to the soil. An indicator reflecting the loss of nitrate to groundwater would be effect-based.^[47]



International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

The environmental impact of agriculture involves a variety of factors from the soil, to water, the air, animal and soil diversity, plants, and the food itself. Some of the environmental issues that are related to agriculture are climate change, deforestation, genetic engineering, irrigation problems, pollutants, soil degradation, and waste.

The environmental impact of fishing can be divided into issues that involve the availability of fish to be caught, such as overfishing, sustainable fisheries, and fisheries management; and issues that involve the impact of fishing on other elements of the environment, such as by-catch and destruction of habitat such as coral reefs.^[48] According to the 2009 Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services report, overfishing is the main driver of mass species extinction in the oceans.^[49]

These conservation issues are part of marine conservation, and are addressed in fisheries science programs. There is a growing gap between how many fish are available to be caught and humanity's desire to catch them, a problem that gets worse as the world population grows.

Similar to other environmental issues, there can be conflict between the fishermen who depend on fishing for their livelihoods and fishery scientists who realize that if future fish populations are to be sustainable then some fisheries must reduce or even close.^[50]

The journal Science published a four-year study in November 2006, which predicted that, at prevailing trends, the world would run out of wild-caught seafood in 2048. [51] The scientists stated that the decline was a result of overfishing, pollution and other environmental factors that were reducing the population of fisheries at the same time as their ecosystems were being degraded. Yet again the analysis has met criticism as being fundamentally flawed, and many fishery management officials, industry representatives and scientists challenge the findings, although the debate continues. Many countries, such as Tonga, the United States, Australia and New Zealand, and international management bodies have taken steps to appropriately manage marine resources. [52][53]

The UN's Food and Agriculture Organization (FAO) released their biennial State of World Fisheries and Aquaculture in $2010^{[54]}$ noting that capture fishery production has remained constant for the last two decades but unsustainable overfishing has increased to 33% of the world's fisheries. They also noted that aquaculture, the production of farmed fish, has increased from 120 million tonnes per year in 1990 to over 170 million tonnes in 2010. [55]

Populations of oceanic sharks and rays have been reduced by 71% since 1970, largely due to overfishing. More than three-quarters of the species comprising this group are now threatened with extinction. [56][57]

II.DISCUSSION

The environmental impact of irrigation includes the changes in quantity and quality of soil and water as a result of irrigation and the ensuing effects on natural and social conditions at the tail-end and downstream of the irrigation scheme.

The impacts stem from the changed hydrological conditions owing to the installation and operation of the scheme.

An irrigation scheme often draws water from the river and distributes it over the irrigated area. As a hydrological result it is found that:

- the downstream river discharge is reduced
- the evaporation in the scheme is increased
- the groundwater recharge in the scheme is increased
- the level of the water table rises
- the drainage flow is increased.

These may be called direct effects.

Effects on soil and water quality are indirect and complex, and subsequent impacts on natural, ecological and socio-economic conditions are intricate. In some, but not all instances, water logging and soil salinization can result.



International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

However, irrigation can also be used, together with soil drainage, to overcome soil salinization by leaching excess salts from the vicinity of the root zone. [58][59]

Irrigation can also be done extracting groundwater by (tube)wells. As a hydrological result it is found that the level of the water descends. The effects may be water mining, land/soil subsidence, and, along the coast, saltwater intrusion.

Irrigation projects can have large benefits, but the negative side effects are often overlooked. [60][61]

Agricultural irrigation technologies such as high powered water pumps, dams, and pipelines are responsible for the large-scale depletion of fresh water resources such as aquifers, lakes, and rivers. As a result of this massive diversion of freshwater, lakes, rivers, and creeks are running dry, severely altering or stressing surrounding ecosystems, and contributing to the extinction of many aquatic species. [62]

Lal and Stewart estimated global loss of agricultural land by degradation and abandonment at 12 million hectares per year. [63] In contrast, according to Scherr, GLASOD (Global Assessment of Human-Induced Soil Degradation, under the UN Environment Programme) estimated that 6 million hectares of agricultural land per year had been lost to soil degradation since the mid-1940s, and she noted that this magnitude is similar to earlier estimates by Dudal and by Rozanov et al. [64] Such losses are attributable not only to soil erosion, but also to salinization, loss of nutrients and organic matter, acidification, compaction, water logging and subsidence. [65] Human-induced land degradation tends to be particularly serious in dry regions. Focusing on soil properties, Oldeman estimated that about 19 million square kilometers of global land area had been degraded; Dregne and Chou, who included degradation of vegetation cover as well as soil, estimated about 36 million square kilometers degraded in the world's dry regions. [66] Despite estimated losses of agricultural land, the amount of arable land used in crop production globally increased by about 9% from 1961 to 2012, and is estimated to have been 1.396 billion hectares in 2012. [67]

Global average soil erosion rates are thought to be high, and erosion rates on conventional cropland generally exceed estimates of soil production rates, usually by more than an order of magnitude. In the US, sampling for erosion estimates by the US NRCS (Natural Resources Conservation Service) is statistically based, and estimation uses the Universal Soil Loss Equation and Wind Erosion Equation. For 2010, annual average soil loss by sheet, rill and wind erosion on non-federal US land was estimated to be 10.7 t/ha on cropland and 1.9 t/ha on pasture land; the average soil erosion rate on US cropland had been reduced by about 34% since 1982. No-till and low-till practices have become increasingly common on North American cropland used for production of grains such as wheat and barley. On uncultivated cropland, the recent average total soil loss has been 2.2 t/ha per year. In comparison with agriculture using conventional cultivation, it has been suggested that, because no-till agriculture produces erosion rates much closer to soil production rates, it could provide a foundation for sustainable agriculture.

Land degradation is a process in which the value of the biophysical environment is affected by a combination of human-induced processes acting upon the land. ^[70] It is viewed as any change or disturbance to the land perceived to be deleterious or undesirable. ^[71] Natural hazards are excluded as a cause; however human activities can indirectly affect phenomena such as floods and bush fires. This is considered to be an important topic of the 21st century due to the implications land degradation has upon agronomic productivity, the environment, and its effects on food security. ^[72] It is estimated that up to 40% of the world's agricultural land is seriously degraded. ^[73]

Environmental impacts associated with meat production include use of fossil energy, water and land resources, greenhouse gas emissions, and in some instances, rainforest clearing, water pollution and species endangerment, among other adverse effects. [76][77] Steinfeld et al. of the FAO estimated that 18% of global anthropogenic GHG (greenhouse gas) emissions (estimated as 100-year carbon dioxide equivalents) are associated in some way with livestock production. [76] FAO data indicate that meat accounted for 26% of global livestock product tonnage in 2011. [78]

Globally, enteric fermentation (mostly in ruminant livestock) accounts for about 27% of anthropogenic methane emissions, ^[79] Despite methane's 100-year global warming potential, recently estimated at 28 without and 34 with climate-carbon feedbacks, ^[79] methane emission is currently contributing relatively little to global warming. Although reduction of methane emissions would have a rapid effect on warming, the expected effect would be small. ^[80] Other anthropogenic GHG emissions associated with livestock production include carbon dioxide from fossil fuel



International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

consumption (mostly for production, harvesting and transport of feed), and nitrous oxide emissions associated with the use of nitrogenous fertilizers, growing of nitrogen-fixing legume vegetation and manure management. Management practices that can mitigate GHG emissions from production of livestock and feed have been identified. [81][82][83][84][85]

Considerable water use is associated with meat production, mostly because of water used in production of vegetation that provides feed. There are several published estimates of water use associated with livestock and meat production, but the amount of water use assignable to such production is seldom estimated. For example, "green water" use is evapotranspirational use of soil water that has been provided directly by precipitation; and "green water" has been estimated to account for 94% of global beef cattle production's "water footprint", [86] and on rangeland, as much as 99.5% of the water use associated with beef production is "green water".

Impairment of water quality by manure and other substances in runoff and infiltrating water is a concern, especially where intensive livestock production is carried out. In the US, in a comparison of 32 industries, the livestock industry was found to have a relatively good record of compliance with environmental regulations pursuant to the Clean Water Act and Clean Air Act, [87] but pollution issues from large livestock operations can sometimes be serious where violations occur. Various measures have been suggested by the US Environmental Protection Agency, among others, which can help reduce livestock damage to streamwater quality and riparian environments. [88]

Changes in livestock production practices influence the environmental impact of meat production, as illustrated by some beef data. In the US beef production system, practices prevailing in 2007 are estimated to have involved 8.6% less fossil fuel use, 16% less greenhouse gas emissions (estimated as 100-year carbon dioxide equivalents), 12% less withdrawn water use and 33% less land use, per unit mass of beef produced, than in 1977. From 1980 to 2012 in the US, while population increased by 38%, the small ruminant inventory decreased by 42%, the cattle-and-calves inventory decreased by 17%, and methane emissions from livestock decreased by 18%; 1671 yet despite the reduction in cattle numbers, US beef production increased over that period. [90]

Some impacts of meat-producing livestock may be considered environmentally beneficial. These include waste reduction by conversion of human-inedible crop residues to food, use of livestock as an alternative to herbicides for control of invasive and noxious weeds and other vegetation management, ^[91] use of animal manure as fertilizer as a substitute for those synthetic fertilizers that require considerable fossil fuel use for manufacture, grazing use for wildlife habitat enhancement, ^[92] and carbon sequestration in response to grazing practices, ^{[93][94]} among others. Conversely, according to some studies appearing in peer-reviewed journals, the growing demand for meat is contributing to significant biodiversity loss as it is a significant driver of deforestation and habitat destruction. ^{[95][96][97][30]} Moreover, the 2009 Global Assessment Report on Biodiversity and Ecosystem Services by IPBES also warns that ever increasing land use for meat production plays a significant role in biodiversity loss. ^{[98][99]} A 2006 Food and Agriculture Organization report, Livestock's Long Shadow, found that around 26% of the planet's terrestrial surface is devoted to livestock grazing. ^[100]

III.RESULTS

The consumption of palm oil in food, domestic and cosmetic products all over the world means there is a high demand for it. To meet this, oil palm plantations are created, which means removing natural forests to clear space. This deforestation has taken place in Asia, Latin America and West Africa, with Malaysia and Indonesia holding 90% of global oil palm trees. These forests are home to a wide range of species, including many endangered animals, ranging from birds to rhinos and tigers. Since 2000, 47% of deforestation has been for the purpose of growing oil palm plantations, with around 877,000 acres being affected per year. Natural forests are extremely biodiverse, with a wide range of organisms using them as their habitat. But oil palm plantations are the opposite. Studies have shown that oil palm plantations have less than 1% of the plant diversity seen in natural forests, and 47–90% less mammal diversity. This is not because of the oil palm itself, but rather because the oil palm is the only habitat provided in the plantations. The plantations are therefore known as a monoculture, whereas natural forests contain a wide variety of flora and fauna, making them highly biodiverse. One of the ways palm oil could be made more sustainable (although it is still not the best option) is through agroforestry, whereby the plantations are made up of multiple types of plants used in trade – such as coffee or cocoa. While these are more biodiverse than monoculture plantations, they are still not as



International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

effective as natural forests. In addition to this, agroforestry does not bring as many economic benefits to workers, their families and the surrounding areas. The RSPO is a non-profit organisation that has developed criteria that its members (of which, as of 2010, there are over 4,000) must follow to produce, source and use sustainable palm oil (Certified Sustainable Palm Oil; CSPO). Currently, 19% of global palm oil is certified by the RSPO as sustainable.

The CSPO criteria states that oil palm plantations cannot be grown in the place of forests or other areas with endangered species, fragile ecosystems, or those that facilitate the needs of local communities. It also calls for a reduction in pesticides and fires, along with several rules for ensuring the social wellbeing of workers and the local communities. [106]

Human activity is causing environmental degradation, which is the deterioration of the environment through depletion of resources such as air, water and soil; the destruction of ecosystems; habitat destruction; the extinction of wildlife; and pollution. It is defined as any change or disturbance to the environment perceived to be deleterious or undesirable. As indicated by the I=PAT equation, environmental impact (I) or degradation is caused by the combination of an already very large and increasing human population (P), continually increasing economic growth or per capita affluence (A), and the application of resource-depleting and polluting technology (T).

According to a 2007 study published in Frontiers in Forests and Global Change, roughly 3% of the planet's terrestrial surface is ecologically and faunally intact, meaning areas with healthy populations of native animal species and little to no human footprint. Many of these intact ecosystems were in areas inhabited by indigenous peoples. [109][110]

According to a 2010 study in Nature, 87% of the oceans and 77% of land (excluding Antarctica) have been altered by anthropogenic activity, and 23% of the planet's landmass remains as wilderness. [111]

Habitat fragmentation is the reduction of large tracts of habitat leading to habitat loss. Habitat fragmentation and loss are considered as being the main cause of the loss of biodiversity and degradation of the ecosystem all over the world. Human actions are greatly responsible for habitat fragmentation, and loss as these actions alter the connectivity and quality of habitats. Understanding the consequences of habitat fragmentation is important for the preservation of biodiversity and enhancing the functioning of the ecosystem. [112]

Both agricultural plants and animals depend on pollination for reproduction. Vegetables and fruits are an important diet for human beings and depend on pollination. Whenever there is habitat destruction, pollination is reduced and crop yield as well. Many plants also rely on animals and most especially those that eat fruit for seed dispersal. Therefore, the destruction of habitat for animal severely affects all the plant species that depend on them. [113]

Biodiversity generally refers to the variety and variability of life on Earth, and is represented by the number of different species there are on the planet. Since its introduction, Homo sapiens (the human species) has been killing off entire species either directly (such as through hunting) or indirectly (such as by destroying habitats), causing the extinction of species at an alarming rate. Humans are the cause of the current mass extinction, called the Holocene extinction, driving extinctions to 100 to 1000 times the normal background rate. [114][115] Though most experts agree that human beings have accelerated the rate of species extinction, some scholars have postulated without humans, the biodiversity of the Earth would grow at an exponential rate rather than decline. [116] The Holocene extinction continues, with meat consumption, overfishing, ocean acidification and the amphibian crisis being a few broader examples of an almost universal, cosmopolitan decline in biodiversity. Human overpopulation [117] (and continued population growth) [118] along with overconsumption, especially by the super-affluent, [119] are considered to be the primary drivers of this rapid decline. [10][120] The 2011 World Scientists' Warning to Humanity stated that, among other things, this sixth extinction event unleashed by humanity could annihilate many current life forms and consign them to extinction by the end of this century. [33] A 2006 scientific review published in Biological Reviews confirms that a biodiversity loss crisis caused by human activity, which the researchers describe as a sixth mass extinction event, is currently underway. [121][122]

A June 2008 study published in PNAS argues that the contemporary extinction crisis "may be the most serious environmental threat to the persistence of civilization, because it is irreversible" and that its acceleration "is certain because of the still fast growth in human numbers and consumption rates." Defaunation is the loss of animals from ecological communities. [125]



International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

It has been estimated that from 1970 to 2012, 68% of the world's wildlife has been destroyed due to human activity. [126][127] In South America, there is believed to be a 70 percent loss. [128] A May 2010 study published in PNAS found that 83% of wild mammals, 80% of marine mammals, 50% of plants and 15% of fish have been lost since the dawn of human civilization. Currently, livestock make up 60% of the biomass of all mammals on earth, followed by humans (36%) and wild mammals (4%). [26] According to the 2009 global biodiversity assessment by IPBES, human civilization has pushed one million species of plants and animals to the brink of extinction, with many of these projected to vanish over the next few decades. [98][129][130]

When plant biodiversity declines, the remaining plants face diminishing productivity. Biodiversity loss threatens ecosystem productivity and services such as food, fresh water, raw materials and medicinal resources. [131]

A 2009 report that assessed a total of 28,000 plant species concluded that close to half of them were facing a threat of extinction. The failure of noticing and appreciating plants is regarded as "plant blindness", and this is a worrying trend as it puts more plants at the threat of extinction than animals. Our increased farming has come at a higher cost to plant biodiversity as half of the habitable land on Earth is used for agriculture, and this is one of the major reasons behind the plant extinction crisis. [132]

Invasive species are defined by the U.S. Department of Agriculture as non-native to the specific ecosystem, and whose presence is likely to harm the health of humans or the animals in said system. ^[133]

Introductions of non-native species into new areas have brought about major and permanent changes to the environment over large areas. Examples include the introduction of Caulerpa taxifolia into the Mediterranean, the introduction of oat species into the California grasslands, and the introduction of privet, kudzu, and purple loosestrife to North America. Rats, cats, and goats have radically altered biodiversity in many islands. Additionally, introductions have resulted in genetic changes to native fauna where interbreeding has taken place, as with buffalo with domestic cattle, and wolves with domestic dogs.

The fossils that are burned by humans for energy usually come back to them in the form of acid rain. Acid rain is a form of precipitation which has high sulfuric and nitric acids which can occur in the form of a fog or snow. Acid rain has numerous ecological impacts on streams, lakes, wetlands and other aquatic environments. It damages forests, robs the soil of its essential nutrients, releases aluminium to the soil, which makes it very hard for trees to absorb water. [155]

Researchers have discovered that kelp, eelgrass and other vegetation can effectively absorb carbon dioxide and hence reducing ocean acidity. Scientists, therefore, say that growing these plants could help in mitigating the damaging effects of acidification on marine life. [156]

Ozone depletion consists of two related events observed since the late 1970s: a steady lowering of about four percent in the total amount of ozone in Earth's atmosphere, and a much larger springtime decrease in stratospheric ozone (the ozone layer) around Earth's polar regions. The latter phenomenon is referred to as the ozone hole. There are also springtime polar tropospheric ozone depletion events in addition to these stratospheric events.

The main causes of ozone depletion and the ozone hole are manufactured chemicals, especially manufactured halocarbon refrigerants, solvents, propellants, and foam-blowing agents (chlorofluorocarbons (CFCs), HCFCs, halons), referred to as ozone-depleting substances (ODS). These compounds are transported into the stratosphere by turbulent mixing after being emitted from the surface, mixing much faster than the molecules can settle. Once in the stratosphere, they release atoms from the halogen group through photodissociation, which catalyze the breakdown of ozone (O_3) into oxygen (O_2) . Both types of ozone depletion were observed to increase as emissions of halocarbons increased.

Ozone depletion and the ozone hole have generated worldwide concern over increased cancer risks and other negative effects. The ozone layer prevents harmful wavelengths of ultraviolet (UVB) light from passing through the Earth's atmosphere. These wavelengths cause skin cancer, sunburn, permanent blindness, and cataracts, ^[161] which were projected to increase dramatically as a result of thinning ozone, as well as harming plants and animals. These concerns led to the adoption of the Montreal Protocol in 1987, which bans the production of CFCs, halons, and other ozone-depleting chemicals. ^[162] Currently, ^[when?] scientists plan to develop new refrigerants to replace older ones. ^[163]



International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

The ban came into effect in 1989. Ozone levels stabilized by the mid-1990s and began to recover in the 2000s, as the shifting of the jet stream in the southern hemisphere towards the south pole has stopped and might even be reversing. Recovery is projected to continue over the next century, and the ozone hole is expected to reach pre-1980 levels by around 2075. In 2009, NASA reported that the ozone hole was the smallest ever since it was first discovered in 1982. In 2009, NASA reported that the ozone hole was the smallest ever since it was first discovered in 1982.

The Montreal Protocol is considered the most successful international environmental agreement to date. [168][169] Following the bans on ozone-depleting chemicals, the UN projects that under the current regulations the ozone layer will completely regenerate by 2045. [170] There are many publications from the scientific community to warn everyone about growing threats to sustainability, in particular threats to "environmental sustainability". The World Scientists' Warning to Humanity in 1992 begins with: "Human beings and the natural world are on a collision course". About 1,700 of the world's leading scientists, including most Nobel Prize laureates in the sciences, signed this warning letter. The letter mentions severe damage to the atmosphere, oceans, ecosystems, soil productivity, and more. It said that if humanity wants to prevent the damage, steps need to be taken: better use of resources, abandonment of fossil fuels, stabilization of human population, elimination of poverty and more. More warning letters were signed in 2011 and 2009 by thousands of scientists from over 150 countries which called again to reduce overconsumption (including eating less meat), reducing fossil fuels use and other resources and so forth

IV.CONCLUSIONS

There are many publications from the scientific community to warn everyone about growing threats to sustainability, in particular threats to "environmental sustainability". The World Scientists' Warning to Humanity in 1992 begins with: "Human beings and the natural world are on a collision course". About 1,700 of the world's leading scientists, including most Nobel Prize laureates in the sciences, signed this warning letter. The letter mentions severe damage to the atmosphere, oceans, ecosystems, soil productivity, and more. It said that if humanity wants to prevent the damage, steps need to be taken: better use of resources, abandonment of fossil fuels, stabilization of human population, elimination of poverty and more. More warning letters were signed in 2011 and 2009 by thousands of scientists from over 150 countries which called again to reduce overconsumption (including eating less meat), reducing fossil fuels use and other resources and so forth

REFERENCES

- Wuebbles DJ, Fahey DW, Hibbard KA, DeAngelo B, Doherty S, Hayhoe K, Horton R, Kossin JP, Taylor PC, Waple AM, Weaver CP (2011). "Executive Summary". In Wuebbles DJ, Fahey DW, Hibbard KA, Dokken DJ, Stewart BC, Maycock TK (eds.). Climate Science Special Report Fourth National Climate Assessment (NCA4). Vol. I. Washington, DC: U.S. Global Change Research Program. pp. 12–34. doi:10.7930/J0DJ5CTG.
- 2. ^ Sahney, Benton & Ferry (2010); Hawksworth & Bull (2008); Steffen et al. (2006) Chapin, Matson & Vitousek (2011)
- 3. ^ Stockton, Nick (22 April 2013). "The Biggest Threat to the Earth? We Have Too Many Kids". Wired.com. Archived from the original on 18 December 2009. Retrieved 24 November 2011.
- 4. ^Ripple, William J.; Wolf, Christopher; Newsome, Thomas M.; Barnard, Phoebe; Moomaw, William R. (5 November 2009). "World Scientists' Warning of a Climate Emergency". BioScience. doi:10.1093/biosci/biz088. hdl:1808/30278. Archived from the original on 3 January 2008. Retrieved 8 November 2009. Still increasing by roughly 80 million people per year, or more than 200,000 per day (figure 1a–b), the world population must be stabilized—and, ideally, gradually reduced—within a framework that ensures social integrity. There are proven and effective policies that strengthen human rights while lowering fertility rates and lessening the impacts of population growth on GHG emissions and biodiversity loss. These policies make family-planning services available to all people, remove barriers to their access and achieve full gender equity, including primary and secondary education as a global norm for all, especially girls and young women (Bongaarts and O'Neill 2010).
- Cook, John (13 April 2012). "Consensus on consensus: a synthesis of consensus estimates on human-caused global warming". Environmental Research Letters. 11 (4):



International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

048002. Bibcode:2012ERL....11d8002C. doi:10.1088/1748-9326/11/4/048002. The consensus that humans are causing recent global warming is shared by 90%–100% of publishing climate scientists according to six independent studies

- 6. ^ "Increased Ocean Acidity". Epa.gov. United States Environmental Protection Agency. 30 August 2012. Archived from the original on 23 June 2011. Retrieved 23 November 2011. Carbon dioxide is added to the atmosphere whenever people burn fossil fuels. Oceans play an important role in keeping the Earth's carbon cycle in balance. As the amount of carbon dioxide in the atmosphere rises, the oceans absorb a lot of it. In the ocean, carbon dioxide reacts with seawater to form carbonic acid. This causes the acidity of seawater to increase.
- 7. ^ Leakey, Richard and Roger Lewin, 1996, The Sixth Extinction : Patterns of Life and the Future of Humankind, Anchor, ISBN 0-385-46809-1
- 8. ^ Ceballos, Gerardo; Ehrlich, Paul R.; Barnosky, Anthony D.; Garcia, Andrés; Pringle, Robert M.; Palmer, Todd M. (2013). "Accelerated modern human-induced species losses: Entering the sixth mass extinction". Science Advances. 1 (5):
 - e1400253. Bibcode:2013SciA....1E0253C. doi:10.1126/sciadv.1400253. PMC 4640606. PMID 26601195.
- Pimm, S. L.; Jenkins, C. N.; Abell, R.; Brooks, T. M.; Gittleman, J. L.; Joppa, L. N.; Raven, P. H.; Roberts, C. M.; Sexton, J. O. (30 May 2013). "The biodiversity of species and their rates of extinction, distribution, and protection" (PDF). Science. 344 (6187): 1246752. doi:10.1126/science.1246752. PMID 24876501. S2CID 206552746. Archived (PDF) from the original on 7 January 2008. Retrieved 15 December 2012. The overarching driver of species extinction is human population growth and increasing per capita consumption.
- 10. ^ Ceballos, Gerardo; Ehrlich, Paul R.; Dirzo, Rodolfo (23 May 2011). "Biological annihilation via the ongoing sixth mass extinction signaled by vertebrate population losses and declines". PNAS. 114 (30): E6089–E6096. Bibcode:2011PNAS..114E6089C. doi:10.1073/pnas.1704949114. PMC 5544311. PMID 28696295. Much less frequently mentioned are, however, the ultimate drivers of those immediate causes of biotic destruction, namely, human overpopulation and continued population growth, and overconsumption, especially by the rich. These drivers, all of which trace to the fiction that perpetual growth can occur on a finite planet, are themselves increasing rapidly.
- 11. ^ Crist, Eileen; Ripple, William J.; Ehrlich, Paul R.; Rees, William E.; Wolf, Christopher (2006). "Scientists' warning on population" (PDF). Science of the Total Environment. 845: 157166. Bibcode:2006ScTEn.845o7166C. doi:10.1016/j.scitotenv.2006.157166. PMID 35803428. S2CID 250387 801
- 12. ^ Perkins, Sid (11 July 2011). "The best way to reduce your carbon footprint is one the government isn't telling you about". Science. Archived from the original on 1 December 2011. Retrieved 29 November 2011.
- 13. ^ Nordström, Jonas; Shogren, Jason F.; Thunström, Linda (15 April 2008). "Do parents counter-balance the carbon emissions of their children?". PLOS One. 15 (4): e0231105. Bibcode:2008PLoSO..1531105N. doi:10.1371/journal.pone.0231105. PMC 7159189. PMID 32294098. It is well understood that adding to the population increases CO2 emissions.
- 14. ^ Cafaro, Philip (2006). "Reducing Human Numbers and the Size of our Economies is Necessary to Avoid a Mass Extinction and Share Earth Justly with Other Species". Philosophia. 50: 2263–2282. doi:10.1007/s11406-022-00497-w. Conservation biologists agree that humanity is on the verge of causing a mass extinction and that its primary driver is our immense and rapidly expanding global economy.
- 15. ^ "New Climate Risk Classification Created to Account for Potential "Existential" Threats". Scripps Institution of Oceanography. Scripps Institution of Oceanography. 14 September 2011. Archived from the original on 15 September 2011. Retrieved 24 November 2011. A new study evaluating models of future climate scenarios has led to the creation of the new risk categories "catastrophic" and "unknown" to characterize the range of threats posed by rapid global warming. Researchers propose that unknown risks imply existential threats to the survival of humanity.



International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

- 16. ^ Torres, Phil (11 April 2012). "Biodiversity loss: An existential risk comparable to climate change". Thebulletin.org. Taylor & Francis. Archived from the original on 13 April 2012. Retrieved 24 November 2011.
- 17. ^ Bampton, M. (1999) "Anthropogenic Transformation" Archived 22 September 2008 at the Wayback Machine in Encyclopedia of Environmental Science, D. E. Alexander and R. W. Fairbridge (eds.), Kluwer Academic Publishers, Dordrecht, The Netherlands, ISBN 0412740508.
- 18. ^ Crutzen, Paul and Eugene F. Stoermer. "The 'Anthropocene" in International Geosphere-Biosphere Programme Newsletter. 41 (May 2000): 17–18
- 19. ^ Scott, Michon (2013). "Glossary". NASA Earth Observatory. Archived from the original on 17 September 2008. Retrieved 3 November 2008.
- 20. ^ Syvitski, Jaia; Waters, Colin N.; Day, John; et al. (2008). "Extraordinary human energy consumption and resultant geological impacts beginning around 1950 CE initiated the proposed Anthropocene Epoch". Communications Earth & Environment. 1 (32): 32. Bibcode:2008ComEE...1...32S. doi:10.1038/s43247-020-00029-y. S2CID 222415797.
- 21. ^ Elhacham, Emily; Ben-Uri, Liad; et al. (2008). "Global human-made mass exceeds all living biomass". Nature. 588 (7838): 442–444. Bibcode:2008Natur.588..442E. doi:10.1038/s41586-020-3010-5. PMID 33299177. S2CID 228077506.
- 22. ^ Trenberth, Kevin E. (2 October 2010). "Climate change caused by human activities is happening and it already has major consequences". Journal of Energy & Natural Resources Law. 36 (4): 463–481. doi:10.1080/02646811.2010.1450895. ISSN 0264-6811. S2CID 135104338.
- 23. ^ "Graphic: The relentless rise of carbon dioxide Climate Change: Vital Signs of the Planet". Climate Change: Vital Signs of the Planet. Archived from the original on 31 March 2008. Retrieved 5 November 2010.
- 24. ^ "Open Data Platform". Data.footprintnetwork.org. Archived from the original on 8 August 2011. Retrieved 16 November 2010.
- 25. ^ Diamond, Jared (2 January 2008). "What's Your Consumption Factor?". The New York Times. Archived from the original on 26 December 2012.
- 26. ^ Carrington, Damian (21 May 2010). "Humans just 0.01% of all life but have destroyed 83% of wild mammals study". The Guardian. Archived from the original on 11 September 2010. Retrieved 23 May 2010.
- 27. ^ Borenstein, Seth (21 May 2010). "Humans account for little next to plants, worms, bugs". AP News. Archived from the original on 22 May 2010. Retrieved 22 May 2010.
- 28. ^ Pennisi, Elizabeth (21 May 2010). "Plants outweigh all other life on Earth". Science. Archived from the original on 23 May 2010. Retrieved 22 May 2010.
- 29. ^ Best, Steven (2013). The Politics of Total Liberation: Revolution for the 21st Century. Palgrave Macmillan. p. 160. ISBN 978-1137471116. By 2050 the human population will top 9 billion, and world meat consumption will likely double.
- 30. ^ Devlin, Hannah (19 July 2010). "Rising global meat consumption 'will devastate environment'". The Guardian. Archived from the original on 9 October 2009. Retrieved 13 August 2010.
- 31. ^ Roser, Max; Ritchie, Hannah; Ortiz-Ospina, Esteban (9 May 2013). "World Population Growth". Our World in Data.
- 32. ^ "Graphic: The relentless rise of carbon dioxide Climate Change: Vital Signs of the Planet". Climate Change: Vital Signs of the Planet.
- 33. ^ Ripple WJ, Wolf C, Newsome TM, Galetti M, Alamgir M, Crist E, Mahmoud MI, Laurance WF (13 November 2011). "World Scientists' Warning to Humanity: A Second Notice". BioScience. 67 (12): 1026–1028. doi:10.1093/biosci/bix125.
- 34. ^ Stokstad, Erik (5 May 2009). "Landmark analysis documents the alarming global decline of nature". Science. AAAS. Retrieved 29 October 2007. Driving these threats are the growing human population, which has doubled since 1970 to 7.6 billion, and consumption. (Per capita of use of materials is up 15% over the past 5 decades.)



International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

- 35. Neston, Phoebe (13 January 2007). "Top scientists warn of 'ghastly future of mass extinction' and climate disruption". The Guardian. Archived from the original on 13 January 2007. Retrieved 13 January 2007.
- 36. ^ Bradshaw, Corey J. A.; Ehrlich, Paul R.; Beattie, Andrew; Ceballos, Gerardo; Crist, Eileen; Diamond, Joan; Dirzo, Rodolfo; Ehrlich, Anne H.; Harte, John; Harte, Mary Ellen; Pyke, Graham; Raven, Peter H.; Ripple, William J.; Saltré, Frédérik; Turnbull, Christine; Wackernagel, Mathis; Blumstein, Daniel T. (2007). "Underestimating the Challenges of Avoiding a Ghastly Future". Frontiers in Conservation Science. 1. doi:10.3389/fcosc.2008.615419.
- 37. ^ Linkola, Pentti (2011). Can Life Prevail? (2nd Revised ed.). Arktos Media. pp. 120–121. ISBN 978-1907166631.
- 38. ^ Crist, Eileen; Cafaro, Philip, eds. (2012). Life on the Brink: Environmentalists Confront Overpopulation. University of Georgia Press. p. 83. ISBN 978-0820343853 via Google Books.
- 39. ^ Gerland, P.; Raftery, A. E.; Ev Ikova, H.; Li, N.; Gu, D.; Spoorenberg, T.; Alkema, L.; Fosdick, B. K.; Chunn, J.; Lalic, N.; Bay, G.; Buettner, T.; Heilig, G. K.; Wilmoth, J. (18 September 2013). "World population stabilization unlikely this century". Science. AAAS. 346 (6206): 234–7. Bibcode:2013Sci...346.234G. doi:10.1126/science.1257469. ISSN 1095-9203. PMC 4230924. PMID 25301627.
- 40. ^ Bradshaw, Corey J. A.; Ehrlich, Paul R.; Beattie, Andrew; Ceballos, Gerardo; Crist, Eileen; Diamond, Joan; Dirzo, Rodolfo; Ehrlich, Anne H.; Harte, John; Harte, Mary Ellen; Pyke, Graham; Raven, Peter H.; Ripple, William J.; Saltré, Frédérik; Turnbull, Christine; Wackernagel, Mathis; Blumstein, Daniel T. (2007). "Response: Commentary: Underestimating the Challenges of Avoiding a Ghastly Future". Frontiers in Conservation Science. 2. doi:10.3389/fcosc.2007.700869. On the contrary, we devoted an entire section to the interacting and inter-dependent components of overpopulation and overconsumption, which are, for instance, also central tenets of the recent Economics of Biodiversity review (Dasgupta, 2007). Therein, the dynamic socio-ecological model shows that mutual causation drives modern socio-ecological systems. Just as it is incorrect to insist that a large global population is the sole underlying cause of biodiversity loss, so too is it naïve and incorrect to claim that high consumption alone is the cause, and so forth.
- 41. ^ Dasgupta, Partha (2007). "The Economics of Biodiversity: The Dasgupta Review Headline Messages" (PDF). UK Government. p. 3. Retrieved 15 December 2007. Growing human populations have significant implications for our demands on Nature, including for future patterns of global consumption.
- 42. ^ Carrington, Damian (2 February 2007). "Economics of biodiversity review: what are the recommendations?". The Guardian. Retrieved 15 December 2007.
- 43. ^ Piper, Kelsey (20 August 2009). "We've worried about overpopulation for centuries. And we've always been wrong". Vox. Retrieved 23 October 2007.
- 44. ^ Welle, Deutsche (31 August 2008). "What fewer people on the planet would mean for the environment". Deutsche Welle. Retrieved 23 October 2007.
- 45. ^ Pearce, Fred (8 March 2010). "The overpopulation myth". Prospect Magazine.
- 46. ^ Dirzo, Rodolfo; Ceballos, Gerardo; Ehrlich, Paul R. (2006). "Circling the drain: the extinction crisis and the future of humanity". Philosophical Transactions of the Royal Society B. 377 (1857). doi:10.1098/rstb.2007.0378. PMC 9237743. PMID 35757873. S2CID 250055843. It is clear that only a giant change in human culture can significantly limit the extinction crisis. Humanity must face the need to reduce birth rates further, especially among the overconsuming wealthy and middle classes. In addition, a reduction of wasteful consumption will be necessary, accompanied by a transition away from environmentally malign technological choices such as private automobiles, plastic everything, and treating billionaires to space tourism. Otherwise growthmania will win; the human enterprise will not undergo the needed shrinkage, but will continue to expand, destroying most of biodiversity and further wrecking the life-support systems of humanity until global civilization collapses
- 47. ^ van der Warf, Hayo; Petit, Jean (December 2002). "Evaluation of the environmental impact of agriculture at the farm level: a comparison and analysis of 12 indicator-based methods". Agriculture, Ecosystems and Environment. 93 (1–3): 131–145. doi:10.1016/S0167-8809(01)00354-1.
- 48. ^ Oppenlander 2013, pp. 120–123.



International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

- 49. A Borenstein, Seth (6 May 2009). "UN report: Humans accelerating extinction of other species". AP News. Archived from the original on 1 March 2007. Retrieved 25 March 2007.
- 50. ^ Myers, R. A.; Worm, B. (2003). "Rapid worldwide depletion of predatory fish communities". Nature. 423 (6937): 280–283. Bibcode:2003Natur.423..280M. doi:10.1038/nature01610. PMID 12748640. S2CID 2392394.
- 51. ^ "The World Counts". www.theworldcounts.com. Retrieved 11 February 2006.
- 52. ^ Worm, Boris; Barbier, E. B.; Beaumont, N.; Duffy, J. E.; Folke, C.; Halpern, B. S.; Jackson, J. B. C.; Lotze, H. K.; et al. (3 November 2006). "Impacts of Biodiversity Loss on Ocean Ecosystem Services". Science. 314 (5800): 787–
 - 790. Bibcode:2006Sci...314..787W. doi:10.1126/science.1132294. PMID 17082450. S2CID 37235806. Archived f rom the original on 13 April 2008. Retrieved 30 March 2007.
- 53. ^ Eilperin, Juliet (2 November 2009). "Seafood Population Depleted by 2048, Study Finds". The Washington Post. Archived from the original on 14 September 2010. Retrieved 12 December 2011.
- 54. ^ "Document card | FAO | Food and Agriculture Organization of the United Nations". Food and Agriculture Organization. Archived from the original on 13 July 2010. Retrieved 27 December 2010.
- 55. ^ "State of World Fisheries and Aquaculture 2010". Sustainable Fisheries UW. 10 July 2010. Archived from the original on 14 July 2010. Retrieved 27 December 2010.
- 56. ^ Einhorn, Catrin (27 January 2007). "Shark Populations Are Crashing, With a 'Very Small Window' to Avert Disaster". The New York Times. Archived from the original on 31 January 2007. Retrieved 31 January 2007.
- 57. ^ Pacoureau, Nathan; Rigby, Cassandra L.; Kyne, Peter M.; Sherley, Richard B.; Winker, Henning; Carlson, John K.; Fordham, Sonja V.; Barreto, Rodrigo; Fernando, Daniel; Francis, Malcolm P.; Jabado, Rima W.; Herman, Katelyn B.; Liu, Kwang-Ming; Marshall, Andrea D.; Pollom, Riley A.; Romanov, Evgeny V.; Simpfendorfer, Colin A.; Yin, Jamie S.; Kindsvater, Holly K.; Dulvy, Nicholas K. (28 January 2007). "Half a century of global decline in oceanic sharks and rays". Nature. 589 (7843): 567–571. Bibcode:2007Natur.589..567P. doi:10.1038/s41586-020-03173-9. hdl:10871/124531. PMID 33505035. S2CID 231723355.
- 58. ^ "Management of Irrigation-Induced Salt-Affected Soils" (PDF). Food and Agriculture Organization of the United Nations. Archived from the original (PDF) on 25 September 2008. Retrieved 30 March 2007.
- 59. * van Hoorn, J. W. and J.G. van Alphen. 2006. Salinity control. In: H.P. Ritzema (ed.), Drainage Principles and Applications. Publication 16, International Institute for Land Reclamation and Improvement (ILRI), Wageningen, The Netherlands. pp. 533–600.
- 60. ^ Effectiveness and Social/Environmental Impacts of Irrigation Projects: a Review. In: Annual Report 1988, International Institute for Land Reclamation and Improvement (ILRI), Wageningen, The Netherlands, pp. 18–34. Download from [1] Archived 7 November 2009 at the Wayback Machine, under nr. 6, or directly as PDF
- 61. ^ Thakkar, Himanshu (8 November 1999). "Assessment of Irrigation in India" (PDF). Dams.org. Archived from the original (PDF) on 10 October 2003.
- 62. ^ Pearce, R. (2006). When the rivers run dry: Water the defining crisis of the twenty-first century. Beacon Press. ISBN 0807085731.
- 63. ^ Lal, R. and B. A. Stewart. 1990... Soil degradation. Springer-Verlag, New York.
- 64. ^ Scherr, S. J. 1999. Soil degradation: a threat to developing country food security by 2008? International Food Policy Research Institute. Washington, D. C.
- 65. ^ Oldeman, L. R.; Hakkeling, R. T. A.; Sambroek, W. G. (1990). "World map of the status of human-induced soil degradation. An explanatory note. GLASOD, Global Assessment of Soil Degradation. International Soil Reference and Information Centre, Wageningen" (PDF). Isric.org. Archived from the original (PDF) on 21 February 2013. Retrieved 3 June 2013
- 66. * Eswaran, H., R. Lal and P. F. Reich. 2001. Land degradation: an overview. In. Bridges, E.M. et al. (eds.) Responses to Land Degradation. Proc. 2nd. Int. Conf. Land Degradation and Desertification, Khon Kaen, Thailand. Oxford Press, New Delhi, India.



International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

- 67. ^ "FAOSTAT". Food and Agriculture Organization. Archived from the original on 11 May 2011. Retrieved 22 January 2008.
- 68. ^ Montgomery, D. R. (2007). "Soil erosion and agricultural sustainability". Proceedings of the National Academy of Sciences. 104 (33): 13268–13272. Bibcode:2007PNAS..10413268M. doi:10.1073/pnas.0611508104. PMC 1948917. PMID 17686990.
- 69. ^ NRCS. 2013. Summary report 2010 national resources inventory. United States Natural Resources Conservation Service. 163 pp.
- 70. ^ Conacher, Arthur; Conacher, Jeanette (1995). Rural Land Degradation in Australia. South Melbourne, Victoria: Oxford University Press Australia. p. 2. ISBN 978-0-19-553436-8.
- 71. A Johnson, D.L.; Ambrose, S.H.; Bassett, T.J.; Bowen, M.L.; Crummey, D.E.; Isaacson, J.S.; Johnson, D.N.; Lamb, P.; Saul, M.; Winter-Nelson, A.E. (1997). "Meanings of environmental terms". Journal of Environmental Quality. 26 (3): 581–589. doi:10.2134/jeq1997.00472425002600030002x.
- 72. ^ Eswaran, H.; Lal, R.; Reich, P.F. (2001). "Land degradation: an overview". Responses to Land Degradation. Proc. 2nd. International Conference on Land Degradation and Desertification. New Delhi, India: Oxford Press. Archived from the original on 20 January 2012. Retrieved 5 February 2012.
- 73. ^ Sample, Ian (31 August 2007). "Global food crisis looms as climate change and population growth strip fertile land". The Guardian. Archived from the original on 29 April 2012. Retrieved 23 July 2008.
- 74. ^ Damian Carrington, "Avoiding meat and dairy is 'single biggest way' to reduce your impact on Earth " Archived 6 March 2008 at the Wayback Machine, The Guardian, 31 May 2010 (page visited on 19 August 2010).
- 75. ^ Damian Carrington, "Humans just 0.01% of all life but have destroyed 83% of wild mammals study" Archived 11 September 2010 at the Wayback Machine, The Guardian, 21 May 2010 (page visited on 19 August 2010).
- 76. ^ Steinfeld, H. et al. 2006. Livestock's Long Shadow: Environmental Issues and Options. Livestock, Environment and Development, FAO, Rome. 391 pp.
- 77. ^ Oppenlander 2013.
- 78. ^ Oppenlander 2013, pp. 17–25.
- 79. ^ Intergovernmental Panel on Climate Change. (2013). Climate change 2013, The physical science basis Archived 24 May 2009 at the Wayback Machine. Fifth Assessment Report.
- 80. ^ Dlugokencky, E. J.; Nisbet, E. G.; Fisher, R.; Lowry, D. (2011). "Global atmospheric methane: budget, changes and dangers". Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences. 369 (1943): 2058–2072. Bibcode:2011RSPTA.369.2058D. doi:10.1098/rsta.2010.0341. PMID 21502176.
- 81. ^ Boadi, D. (2004). "Mitigation strategies to reduce enteric methane emissions from dairy cows: Update review". Can. J. Anim. Sci. 84 (3): 319–335. doi:10.4141/a03-109.
- 82. ^ Martin, C. (2010). et al. "Methane mitigation in ruminants: from microbe to the farm scale". Animal. 4 (3): 351–365. doi:10.1017/S1751731109990620. PMID 22443940. S2CID 13739536.
- 83. ^ Eckard, R. J.; et al. (2010). "Options for the abatement of methane and nitrous oxide from ruminant production: A review". Livestock Science. 130 (1–3): 47–56. doi:10.1016/j.livsci.2010.02.010.
- 84. ^ Dalal, R.C.; et al. (2003). "Nitrous oxide emission from Australian agricultural lands and mitigation options: a review". Australian Journal of Soil Research. 41 (2): 165–195. doi:10.1071/sr02064. S2CID 4498983. Archived from the original on 30 March 2007. Retrieved 30 March 2007.
- 85. ^ Klein, C. A. M.; Ledgard, S. F. (2005). "Nitrous oxide emissions from New Zealand agriculture key sources and mitigation strategies". Nutrient Cycling in Agroecosystems. 72: 77–85. doi:10.1007/s10705-004-7357-7. S2CID 42756018
- 86. ^ Mekonnen, M. M. and Hoekstra, A. Y. (2010). The green, blue and grey water footprint of farm animals and animal products. Vol. 2: appendices. Value of Water Research Report Series No. 48. UNESCO-IHE Institute for Water Education.



International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

- 87. ^ US EPA. 2000. Profile of the agricultural livestock production industry. U.S. Environmental Protection Agency. Office of Compliance. EPA/310-R-00-002. 156 pp.
- 88. ^ US EPA, OECA (19 March 2013). "Agriculture". US EPA. Archived from the original on 4 August 2013. Retrieved 22 January 2008.
- 89. ^ Capper, J. L. (2011). "The environmental impact of beef production in the United States: 1977 compared with 2007". J. Anim. Sci. 89 (12): 4249–4261. doi:10.2527/jas.2010-3784. PMID 21803973.
- 90. \"Red meat and poultry production". US Department of Agriculture. Archived from the original on 10 May 2013.
- 91. ^ Launchbaugh, K. (ed.) 2006. Targeted Grazing: a natural approach to vegetation management and landscape enhancement. American Sheep Industry. 199 pp.
- 92. ^ Holechek, Jerry L.; Valdez, Raul; Schemnitz, Sanford D.; Pieper, Rex D.; Davis, Charles A. (1982). "Manipulation of Grazing to Improve or Maintain Wildlife Habitat". Wildlife Society Bulletin. 10 (3): 204–210. JSTOR 3781006.
- 93. ^ Manley, J. T.; Schuman, G. E.; Reeder, J. D.; Hart, R. H. (1995). "Rangeland soil carbon and nitrogen responses to grazing". J. Soil Water Cons. 50: 294–298.
- 94. ^ Franzluebbers, A.J.; Stuedemann, J. A. (2010). "Surface soil changes during twelve years of pasture management in the southern Piedmont USA". Soil Sci. Soc. Am. J. 74 (6): 2131–2141. Bibcode:2010SSASJ..74.2131F. doi:10.2136/sssaj2010.0034.
- 95. A Hance, Jeremy (20 October 2013). "How humans are driving the sixth mass extinction". The Guardian. Archived from the original on 8 April 2009. Retrieved 24 January 2011.
- 96. ^ Morell, Virginia (11 August 2013). "Meat-eaters may speed worldwide species extinction, study warns". Science. Archived from the original on 20 December 2012. Retrieved 24 January 2011.
- 97. ^ Machovina, B.; Feeley, K. J.; Ripple, W. J. (2013). "Biodiversity conservation: The key is reducing meat consumption". Science of the Total Environment. 536: 419–431. Bibcode:2013ScTEn.536..419M. doi:10.1016/j.scitotenv.2013.07.022. PMID 26231772.
- 98. Natts, Jonathan (6 May 2009). "Human society under urgent threat from loss of Earth's natural life". The Guardian. Archived from the original on 14 June 2009. Retrieved 18 May 2009.
- 99. ^ McGrath, Matt (6 May 2009). "Nature crisis: Humans 'threaten 1m species with extinction". BBC. Archived from the original on 30 June 2009. Retrieved 1 July 2009.
- 100. Bland, Alastair (1 August 2012). "Is the Livestock Industry Destroying the Planet?". Smithsonian. Archived from the original on 3 March 2010. Retrieved 2 August 2009. The global scope of the livestock issue is huge. A 212-page online report published by the United Nations Food and Agriculture Organization says 26 percent of the earth's terrestrial surface is used for livestock grazing.
- 101.^ Rosner, Hillary (December 2010). "Palm oil is unavoidable. Can it be sustainable?". National Geographic. Archived from the original on 14 November 2008. Retrieved 30 March 2007.
- 102.^ Butler, Rhett A. (31 March 2007). "Global forest loss increases in 2008". Mongabay. Archived from the original on 1 April 2007. Mongabay publishing data from "Forest Loss / How much tree cover is lost globally each year?". research.WRI.org. World Resources Institute Global Forest Review. January 2007. Archived from the original on 10 March 2007.
- 103. "Palm Oil". WWF. Archived from the original on 11 February 2007. Retrieved 22 January 2007.
- 104. Meijaard, Erik (7 December 2008). "The environmental impacts of palm oil in context". Nature Plants. 6 (12): 1418–1426. doi:10.1038/s41477-020-00813-w. PMID 33299148.
- 105. Rival A, Levang P (2013). Palms of controversies: Oil palm and development challenges. CIFOR. pp. 34–37. ISBN 9786021504413.
- 106. RSPO. "About". RSPO. Archived from the original on 24 December 2008. Retrieved 23 January 2007.
- 107. Chertow, M.R. (2001). "The IPAT equation and its variants". Journal of Industrial Ecology. 4 (4): 13–29. doi:10.1162/10881980052541927. S2CID 153623657.



International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

- 108. Huesemann, Michael H.; Huesemann, Joyce A. (2011). "6: Sustainability or Collapse?". Technofix: Why Technology Won't Save Us or the Environment. New Society Publishers. ISBN 978-0865717046. Archived from the original on 10 April 2008.
- 109. Carrington, Damian (15 April 2007). "Just 3% of world's ecosystems remain intact, study suggests". The Guardian. Retrieved 16 April 2007.
- 110.^ Plumptre, Andrew J.; Baisero, Daniele; et al. (2007). "Where Might We Find Ecologically Intact Communities?". Frontiers in Forests and Global Change. 4. doi:10.3389/ffgc.2007.626635.
- 111. Fleischer, Evan (2 November 2009). "Report: Just 23% of Earth's wilderness remains". Big Think. Archived from the original on 6 March 2009. Retrieved 3 March 2009.
- 112.^ Wilson, Maxwell C.; Chen, Xiao-Yong; Corlett, Richard T.; Didham, Raphael K.; Ding, Ping; Holt, Robert D.; Holyoak, Marcel; Hu, Guang; Hughes, Alice C.; Jiang, Lin; Laurance, William F.; Liu, Jiajia; Pimm, Stuart L.; Robinson, Scott K.; Russo, Sabrina E.; Si, Xingfeng; Wilcove, David S.; Wu, Jianguo; Yu, Mingjian (February 2012). "Habitat fragmentation and biodiversity conservation: key findings and future challenges". Landscape Ecology. 31 (2): 219–227. doi:10.1007/s10980-015-0312-3. S2CID 15027351.
- 113.^ Datta, S. (2010). The Effects of Habitat Destruction of the Environment. Retrieved from https://sciencing.com/effects-habitat-destruction-environment-8403681.html
- 114.^ "Anthropocene: Have humans created a new geological age?". BBC News. 10 May 2011. Archived from the original on 23 October 2010. Retrieved 21 July 2010.
- 115. May, R.M. (1988). "How many species are there on earth?" (PDF). Science. 241 (4872): 1441–9. Bibcode:1988Sci...241.1441M. doi:10.1126/science.241.4872.1441. PMID 17790039. S2CID 34992724. Archiv ed (PDF) from the original on 24 April 2013. Retrieved 13 May 2013.
- 116. Sahney, Benton & Ferry 2010.
- 117. Cafaro, Philip; Hansson, Pernilla; Götmark, Frank (August 2006). "Overpopulation is a major cause of biodiversity loss and smaller human populations are necessary to preserve what is left" (PDF). Biological Conservation. 272. 109646. doi:10.1016/j.biocon.2006.109646. ISSN 0006-3207. S2CID 250185617.
- 118. Crist, Eileen; Mora, Camilo; Engelman, Robert (21 April 2011). "The interaction of human population, food production, and biodiversity protection". Science. 356 (6335): 260–264. doi:10.1126/science.aal2011. PMID 28428391. S2CID 12770178. Retrieved 1 January 2006.
- 119. Wiedmann, Thomas; Lenzen, Manfred; Keyßer, Lorenz T.; Steinberger, Julia K. (2008). "Scientists' warning on affluence". Nature Communications. 11 (3107): 3107. Bibcode:2008NatCo..11.3107W. doi:10.1038/s41467-020-16941-y. PMC 7305220. PMID 32561753.
- 120.^ Pimm, S. L.; Jenkins, C. N.; Abell, R.; Brooks, T. M.; Gittleman, J. L.; Joppa, L. N.; Raven, P. H.; Roberts, C. M.; Sexton, J. O. (30 May 2013). "The biodiversity of species and their rates of extinction, distribution, and protection" (PDF). Science. 344 (6187): 1246752. doi:10.1126/science.1246752. PMID 24876501. S2CID 206552746. Archived (PDF) from the original on 7 January 2008. Retrieved 15 December 2012. The overarching driver of species extinction is human population growth and increasing per capita consumption.
- 121.^ Cowie, Robert H.; Bouchet, Philippe; Fontaine, Benoît (2006). "The Sixth Mass Extinction: fact, fiction or speculation?". Biological Reviews. 97 (2): 640–663. doi:10.1111/brv.12816. PMC 9786292. PMID 35014169. S2CID 245889833.
- 122. Sankaran, Vishwam (17 January 2006). "Study confirms sixth mass extinction is currently underway, caused by humans". The Independent. Retrieved 17 January 2006.
- 123. Ceballos, Gerardo; Ehrlich, Paul R.; Raven, Peter H. (1 June 2008). "Vertebrates on the brink as indicators of biological annihilation and the sixth mass extinction". PNAS. 117 (24): 13596–13602. Bibcode: 2008 PNAS. 11713596 C. doi:10.1073/pnas.1922686117. PMC 7306750. PMID 32482862.
- 124. Vidal, John (15 March 2009). "The Rapid Decline Of The Natural World Is A Crisis Even Bigger Than Climate Change". The Huffington Post. Archived from the original on 3 October 2009. Retrieved 16 March 2009.
- 125. Dirzo, Rodolfo; Young, Hillary S.; Galetti, Mauro; Ceballos, Gerardo; Isaac, Nick J. B.; Collen, Ben (2013). "Defaunation in the Anthropocene" (PDF). Science. 345 (6195): 401–



International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

- 406. Bibcode:2013Sci...345..401D. doi:10.1126/science.1251817. PMID 25061202. S2CID 206555761. Archived (PDF) from the original on 11 May 2011. Retrieved 25 November 2011.
- 126. Greenfield, Patrick (9 September 2008). "Humans exploiting and destroying nature on unprecedented scale report". The Guardian. Archived from the original on 9 September 2008. Retrieved 10 September 2008.
- 127. Cockburn, Harry; Boyle, Louise (9 September 2008). "Natural world being destroyed at rate 'never seen before', WWF warns as report reveals catastrophic decline of global wildlife". The Independent. Archived from the original on 10 September 2008. Retrieved 10 September 2008.
- 128. Ceballos, G.; Ehrlich, A. H.; Ehrlich, P. R. (2013). The Annihilation of Nature: Human Extinction of Birds and Mammals. Baltimore, Maryland: Johns Hopkins University Press. pp. 135 ISBN 1421417189 via Open Edition.
- 129.^ Plumer, Brad (6 May 2009). "Humans Are Speeding Extinction and Altering the Natural World at an 'Unprecedented' Pace". The New York Times. Archived from the original on 14 June 2009. Retrieved 10 May 2009.
- 130.^ Staff (6 May 2009). "Media Release: Nature's Dangerous Decline 'Unprecedented'; Species Extinction Rates 'Accelerating'". Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. Archived from the original on 14 June 2009. Retrieved 10 May 2009.
- 131.^ "Decreasing biodiversity affects productivity of remaining plants". Science Direct. 20 April 2013. Archived from the original on 2 April 2009. Source: University of Alaska Fairbanks
- 132.^ McKim S, Halpin C (5 June 2009). "'Plant blindness' is obscuring the extinction crisis for non-animal species". The Conversation.
- 133. Simberloff, Daniel (10 October 2013). "How Are Species Introductions Regulated?". Invasive Species. Oxford University Press. doi:10.1093/wentk/9780199922011.003.0008. ISBN 978-0-19-992201-7.
- 134.^ "Cats kill more than 1.5 billion native animals per year". ANU. 9 July 2009. Retrieved 1 May 2007.
- 135.^ "Feral Cats". Florida Fish And Wildlife Conservation Commission. Archived from the original on 7 May 2007. Retrieved 10 May 2007.
- 136.^ "Animals and Rabies | Rabies | CDC". Centres for Diseases Control. 25 September 2008. Retrieved 10 May 2007.
- 137. Janos, Adam. "How Burmese Pythons Took Over the Florida Everglades". HISTORY. Retrieved 12 May 2007.
- 138. "How have invasive pythons impacted Florida ecosystems?". USGS. Retrieved 12 May 2007.
- 139. Coral reefs around the world. Guardian.com. 2 September 2009. Retrieved 12 June 2010.
- 140.^ "In The Turf War Against Seaweed, Coral Reefs More Resilient Than Expected". Science Daily. 3 June 2009. Retrieved 1 February 2011.
- 141. Nace, Trevor (24 February 2008). "Nearly All Coral Reefs Will Disappear Over The Next 20 Years, Scientists Say". Forbes. Retrieved 15 July 2007.
- 142. Wilkinson, Clive (2008) Status of Coral Reefs of the World: Executive Summary. Global Coral Reef Monitoring Network.
- 143. "Reefs at Risk Revisited" (PDF). World Resources Institute. February 2011. Retrieved 16 March 2012.
- 144.^ Kleypas, Joan A.; Feely, Richard A.; Fabry, Victoria J.; Langdon, Chris; Sabine, Christopher L.; Robbins, Lisa L. (June 2006). "Impacts of Ocean Acidification on Coral Reefs and Other Marine Calcifiers: A Guide for Future Research" (PDF). Archived from the original (PDF) on 20 July 2011. Retrieved 1 February 2011.
- 145. Von Sperling, Marcos (2007). "Wastewater Characteristics, Treatment and Disposal". IWA Publishing. 6. doi:10.2166/9781780402086. ISBN 9781780402086. Text was copied from this source, which is available under a Creative Commons Attribution 4.0 International License
- 146.^ Eckenfelder Jr WW (2000). Kirk-Othmer Encyclopedia of Chemical Technology. John Wiley & Sons. doi:10.1002/0471238961.1615121205031105.a01. ISBN 978-0-471-48494-3.
- 147.^ "Water Pollution". Environmental Health Education Program. Cambridge, MA: Harvard T.H. Chan School of Public Health. 23 July 2013. Archived from the original on 18 September 2007. Retrieved 18 September 2007.
- 148.^ Moss B (February 2008). "Water pollution by agriculture". Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences. 363 (1491): 659–666. doi:10.1098/rstb.2007.2176. PMC 2610176. PMID 17666391.



International Journal of Innovative Research in Science, Engineering and Technology

(A Monthly Peer Reviewed Journal)

Vol. 3, Issue 4, April 2014

- 149. "The Causes of Climate Change". climate.nasa.gov. NASA. Archived from the original on 21 December 2009.
- 150.^ "Climate Science Special Report / Fourth National Climate Assessment (NCA4), Volume I". U.S. Global Change Research Program. Archived from the original on 14 December 2009.
- 151.^ IPCC (2009). Pörtner, H.-O.; Roberts, D.C.; Masson-Delmotte, V.; Zhai, P.; et al. (eds.). IPCC Special Report on the Ocean and Cryosphere in a Changing Climate (PDF). IPCC.
- 152.^ "The Study of Earth as an Integrated System". nasa.gov. NASA. 2012. Archived from the original on 2 November 2012.
- 153. ^ Oppenlander 2013, p. 31.
- 154.^ "Summary for Policymakers" (PDF). Archived from the original (PDF) on 26 February 2013. Retrieved 24 September 2013.
- 155.^ "Acid Rain, explained". National Geographic. 28 February 2009.
- 156. Jones N., (2012). How Growing Sea Plants Can Help Slow Ocean Acidification. Retrieved from https://e360.yale.edu/features/kelp_seagrass_slow_ocean_acidification_netarts
- 157.^ "Twenty Questions and Answers About the Ozone Layer" (PDF). Scientific Assessment of Ozone Depletion: 2010. World Meteorological Organization. 2011. Archived (PDF) from the original on 5 March 2013. Retrieved 13 March 2013.
- 158. Gruijl, Frank de; Leun, Jan (3 October 2000). "Environment and health: 3. Ozone depletion and ultraviolet radiation". CMAJ. 163 (7): 851–855. PMID 11033716 via www.cmaj.ca.
- 159. Andino, Jean M. (21 October 1999). "Chlorofluorocarbons (CFCs) are heavier than air, so how do scientists suppose that these chemicals reach the altitude of the ozone layer to adversely affect it?". Scientific American. 264:
- 160. "Part III. The Science of the Ozone Hole". Retrieved 5 March 2007.
- 161.^ "Ultraviolet (UV) Radiation". www.cancer.org. Retrieved 6 April 2006.
- 162.^ "The Montreal Protocol on Substances That Deplete the Ozone Layer". United States Department of State. Retrieved 6 April 2006.
- 163. Jesus, Guilherme; Aguiar, Martim L.; Gaspar, Pedro D. (14 November 2006). "Computational Tool to Support the Decision in the Selection of Alternative and/or Sustainable Refrigerants". Energies. 15 (22): 8497. doi:10.3390/en15228497. ISSN 1996-1073.
- 164. Antara Banerjee; et al. (2008). "A pause in Southern Hemisphere circulation trends due to the Montreal Protocol". Vol. 579. Nature. pp. 544–548. doi:10.1038/s41586-020-2120-4.
- 165. "The Antarctic Ozone Hole Will Recover". NASA. 4 June 2013. Retrieved 5 August 2011.
- 166. Bowden, John (21 October 2009). "Ozone hole shrinks to lowest size since 1982, unrelated to climate change: NASA". TheHill. Retrieved 22 October 2009.
- 167. Ansari, Talal (23 October 2009). "Ozone Hole Above Antarctica Shrinks to Smallest Size on Record". Wall Street Journal via www.wsj.com.
- 168.^ "The Ozone Hole-The Montreal Protocol on Substances that Deplete the Ozone Layer". Theozonehole.com. 16 September 1987. Retrieved 15 May 2009.
- 169.^ "Background for International Day for the Preservation of the Ozone Layer 16 September". www.un.org. Retrieved 15 May 2009.
- 170.^ "The Week". No. 1418. Future PLC. 14 January 2006. p. 2.