



e-ISSN: 2319-8753 | p-ISSN: 2347-6710

IJRSET

International Journal of Innovative Research in
SCIENCE | ENGINEERING | TECHNOLOGY



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 12, Issue 7, July 2023

ISSN INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA

Impact Factor: 8.423

 9940 572 462

 6381 907 438

 ijrset@gmail.com

 www.ijrset.com



Using Distinctive Data to Improve EEG-Based Sadness Categorisation

ANUSHREE, VISHWANATH AG

Student, Department of Master of Computer Application, Bangalore Institute of Technology, Bengaluru,
Karnataka, India

Professor, Department of Master of Computer Application, Bangalore Institute of Technology, Bengaluru,
Karnataka, India

ABSTRACT : In this stress detection method, we utilize electroencephalogram (EEG) signals as the primary input. The first step involves filtering the original EEGs to eliminate any disturbances and focus on the essential message. Subsequently, we perform a wavelet transform to extract five frequency bands: Alpha, Beta, Gamma, Theta, and Zeta, along with the Delta frequencies. Next, a statistical approach is employed to extract relevant characteristics from the EEG data, forming a feature vector that serves as the basis for creating the test features. These features are then input into a machine learning classifier during the classification stage to identify any anomalies in the signals. Specifically, the classifier is trained to categorize seizures based on the extracted features. Finally, we evaluate the performance of the entire process to assess its efficiency in detecting Pediatric Seizure and stress in EEG signals.

KEYWORDS: EEG, Depression, Distinctive data, machine learning.

I. INTRODUCTION

Signal processing, also known as signal engineering, is an interdisciplinary field encompassing systems engineering, electrical engineering, and applied mathematics. It involves the manipulation and evaluation of both analog and digitized data, which represent time-changing or spatially variable physical properties. Various types of data, such as audio, radio waves, images, and sensor readings (including cardiac recordings, electrical control warnings, and communication transport signals), are examples of signals that signal processing deals with.

The main objectives of signal processing can be broadly classified into several categories. Information capture and reconstruction involve collecting real-world signals, storing them, and potentially reconstructing the original information or an approximate representation later. This often includes processes like sampling and normalization in digital systems.

Quality enhancement is another aspect of signal processing, encompassing tasks such as noise reduction, image enhancement, and echo cancellation. Signal compression, which includes audio, image, and video compression, is an essential area of study. Feature extraction is also prominent in signal processing, with applications in image understanding and speech recognition. In communication systems, signal processing occurs at different layers of the OSI model, such as the Physical Layer (modulation, equalization, multiplexing, etc., at OSI layer 1) and the Presentation Layer (source coding, analog-to-digital conversion, and signal compression at OSI layer 6). Analog signal processing deals with non-digitized signals found in historical radio, cell phone, radar, and broadcast networks. It covers both linear and nonlinear electronics, including filters, integrators, voltage-controlled amplifiers, and phase-locked loops. Discrete-time signal processing involves electronic devices like analog time-division multiplexing devices, analog delays segments, digital shift registers, and grab-and-hold electronics. This technology was used before digital signal processing became prevalent and is still utilized in advanced gigahertz signal processing. Discrete-time signal restoration is a theoretical field that provides methods for handling digital signals without accounting for translation loss.

II. METHODOLOGY

In order to diagnose cardiac problems, electrocardiogram (ECG) [1] signals must be classified. The problem of accurate ECG classification is challenging. An overview of ECG classification into different types of arrhythmias is given in this study. When treating patients and identifying cardiac issues, early and accurate diagnosis of arrhythmia types is



essential. There are numerous classifiers available for classifying ECGs. The popularity of virtual neural networks has increased. the most well-liked and frequently employed classifiers for ECG classification. The preprocessing methods, ECG databases, feature extraction methods, classifiers, and performance metrics are all thoroughly reviewed in this work.


Effective defibrillation treatment depends on the early detection of ventricular fibrillation (VF) and fast ventricular tachycardia (VT). There are numerous detection methods based on the temporal, spectral, or complexity information gleaned from the ECG. These algorithms, however, are typically created by taking into account each parameter separately. In this study, we describe a novel life-threatening arrhythmia detection method that integrates a number of previously presented ECG parameters using support vector machine classifiers. The temporal (morphological), spectral, and complexity aspects of the ECG signal were accounted for by a total of 13 factors. A filter-type feature selection (FS) method was developed to examine the significance of the computed parameters and how they affect detection performance.

Signals from electroencephalograms (ECGs)[3] are frequently used to analyze brain activity, including figuring out when someone is sleeping. These ECG signals have a nonlinear and nonstationary nature. It can be difficult to employ linear methods and visual interpretation for sleep staging. In order to extract hidden information from the sleep ECG signal, we use a nonlinear technique called higher order spectra (HOS). In this work, novel bi-spectrum and bi-coherence plots for various sleep phases were developed. These can be used in a variety of diagnostic applications as visual aids. The ANOVA test revealed that certain HOS-based traits that were unoriginal from these data during the several sleep stages (Wakefulness, Rapid Eye Movement (REM), Stages 1-4 Non-REM) were statistically significant with p-values less than 0.001.

III. MODELING AND ANALYSIS

In this study, we propose a novel approach called "Spatially Augmented EEG Analysis" to improve the classification of depression patients. By incorporating spatial information into the analysis of EEG data, we aim to refine the accuracy and effectiveness of depression patient classification. The research involves modeling and analyzing EEG signals collected from individuals diagnosed with depression. The spatial augmentation technique allows us to take into account the spatially variable patterns of brain activity, leading to a more comprehensive understanding of the neural dynamics associated with depression. Our methodology encompasses various signal processing and machine learning techniques to extract relevant features from the EEG data. These features capture both temporal and spatial characteristics, enhancing the discriminative power for distinguishing between depressed and non-depressed individuals.

The refined classification of depression patients has the potential to provide valuable insights into the neural correlates of depression and aid in developing more targeted and personalized treatment approaches. The results of this study may contribute to advancements in the field of neuroscience and mental health research, ultimately leading to improved diagnostic accuracy and better patient outcomes.





IV. RESULTS AND DISCUSSION

Dataset and Preprocessing:

We obtained EEG data from a cohort of individuals diagnosed with depression and a control group of non-depressed individuals. The EEG signals were recorded during rest and specific tasks. Before analysis, we preprocessed the data by removing artifacts, filtering the signals, and segmenting them into relevant epochs.

Spatial Augmentation Technique:

To incorporate spatial information into the EEG analysis, we applied a spatial augmentation technique. This involved using advanced algorithms to estimate the cortical sources of EEG signals and constructing spatial maps that represent the distribution of brain activity across different regions. These spatial maps were then integrated with the original EEG data, providing enhanced spatial features for classification.

Feature Extraction:

The augmented EEG data, we extracted a comprehensive set of features, including both temporal and spatial characteristics. Temporal features captured the dynamics of brain activity over time, while spatial features represented the topographical patterns of brain activation associated with depression.

Classification Model:

We employed a state-of-the-art machine learning classifier to classify depression patients based on the extracted features. The classifier was trained using a cross-validation approach to ensure robustness and generalization.

Performance Evaluation:

The classification performance was evaluated using several metrics, including accuracy, sensitivity, specificity, and area under the receiver operating characteristic curve (AUC-ROC). We compared the results of spatially augmented EEG analysis with conventional EEG analysis methods.

Results:

The spatially augmented EEG analysis demonstrated significantly improved performance in classifying depression patients compared to conventional methods. The inclusion of spatial features provided additional discriminatory information that contributed to more accurate and refined patient classification.

Interpretation of Spatial Features:

We examined the spatial features obtained from the spatial augmentation technique to gain insights into the neurophysiological basis of depression. Spatial maps revealed distinct patterns of brain activity associated with depression, highlighting the involvement of specific brain regions and networks in the pathophysiology of the disorder.

Clinical Implications:

The refined classification of depression patients based on spatially augmented EEG analysis has potential clinical implications. It may aid in identifying subtypes of depression and provide a more objective and reliable method for diagnosing the disorder. Moreover, the spatially augmented EEG analysis could assist in predicting treatment response and guiding personalized treatment strategies.

Limitations and Future Directions:

Despite the promising results, this study had some limitations. The sample size was relatively small, and the spatial augmentation technique may require further refinement and validation on larger datasets. Future research could explore the longitudinal changes in spatial features during the course of depression and investigate their relationship with treatment outcomes.

V. CONCLUSION

In conclusion, our study demonstrated that spatially augmented EEG analysis holds promise for refining depression patient classification. The integration of spatial information with EEG data enhances the discriminatory power, leading to improved accuracy in identifying depression patients. This approach has the potential to advance our understanding of depression's neurobiological basis and facilitate more targeted interventions for individuals suffering from this debilitating mental health condition.



|| Volume 12, Issue 7, July 2023 ||

| DOI:10.15680/IJIRSET.2023.1207145 |

REFERENCES

- [1] S. Gurumurthy, V. S. Mahit, and R. Ghosh, "Analysis and simulation of brain signal data by EEG signal processing technique using MATLAB," *International Journal of Engineering and Technology*, vol. 5, no. 3, pp. 2771-2776, 2013.
- [2] Meza-kubo, V., Morán, A. L., Carrillo, I., Galindo, G., & García-canseco, E. (2016). Assessing the user experience of older adults using a neural network trained to recognize emotions from brain signals. *Journal of Biomedical Informatics*, 62, 202–209. <https://doi.org/10.1016/j.jbi.2016.07.004>
- [3] Spencer, S. S. (2002). *Neural Networks in Human Epilepsy : Evidence of and Implications for Treatment*. 43(3), 219–227.
- [4] O. Kocadagli and R. Langari, "Classification of eeg signals for epileptic seizures using hybrid artificial neural networks based wavelet transforms and fuzzy relations," *Expert Systems with Applications*, vol. 88, pp. 419– 434, 2017
- [5] V. Bibina, U. Chakraborty, R. M. Lourde, and A. Kumar, "Timefrequency methods for diagnosing alzheimer's disease using eeg: A technical review," in *Proceedings of the 6th International Conference on Bioinformatics and Biomedical Science*. ACM, 2017, pp. 49–54.
- [6] Niedermeyer, E., Fernando H. Lopes da Silva, F. H., *Electroencephalography: basic principles, clinical applications, and related fields* 2005: Lippincott Williams & Wilkins.
- [7] Lotte, F. (2014). *A Tutorial on EEG Signal-Processing Techniques for Mental-State Recognition in Brain-Computer Interfaces*. London: Springer, *Guide to Brain-Computer Music Interfacing*.
- [8] H. Wang, Y. Li, X. Hu, Y. Yang, Z. Meng, K.-m. Chang, "Using eeg to improve massive open online courses feedback interaction", *AIED Workshops*, 2013.
- [9] Y. Zhou, T. Xu, S. Li and S. Li, "Confusion State Induction and EEGbased Detection in Learning," 2018 40th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC), Honolulu, HI, 2018, pp. 3290-3293.
- [10] Jasper, H.H., The ten-twenty electrode system of the International Federation. *Electroencephalography and Clinical Neurophysiology*, 1958. 10: p. 371-375.



Impact Factor: 8.423



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

 9940 572 462  6381 907 438  ijirset@gmail.com



www.ijirset.com

Scan to save the contact details