

Volume 9, Issue 9, September 2020



Impact Factor: 7.512





|| Volume 9, Issue 9, September 2020 ||

Epidemiological Study on Head & Neck Cancer in Various Districts of Nepal 2013 – 2015

Prativa Neupane ^{1*,} Dej Kumar Gautam ¹, Benju Rashmi Pradhan ¹, Suk Bahadur Tamang ¹,

Khem Bahadur Karki², Krishna Prasad Subedi¹

B.P.Koirala Memorial Cancer Hospital, Bharatpur, Nepal¹ Institute of Medicine, Tribhuvan University, Nepal²

*Corresponding Author

ABSTRACT: Head and neck cancer is a group of cancers that starts in the mouth, nose, throat, larynx, sinuses, or salivary glands etc.

In Nepal total 9718 new cancer cases were reported by hospital based national cancer registration programme. Whereas, 1882 (19.3%) were head & neck cancer, which was only 1363 (19.2%) out of 8729 cases in 2011 and 1401 (19.4%) in 2012. Similarly, 1741 (19.9%) and 1868 (20.3%) cases of oral cancer were diagnosed for 2013 and 2014.

It shows cancer Head & neck cancer in Nepal was increasing every year. Exact cancer incidence, death and relevant community based details of cancer in Nepal is unknown to address the need of community based epidemiological study was initiated community based cancer research in 15 districts of four state of Nepal. In this study we focus to determine the current status and trend of head & neck cancer.

This was a descriptive epidemiological study with primary and secondary data analysis of new oral cancer that are recorded and collected in different data source institution at 15 districts of four state of Nepal science 1st January to 31st December 2013,2014 & 2015. The data analysis was carried out by using SPSS 19.0 Head & neck cancer was relatively increased for 2013 to 2014. Whereas, releatively decreased in 2015.

The trend of head & neck cancer was seen to be fluctuating yearly during the study period. Though well organized awareness activities, and screening programs are still needed to prevent and control the incidence of head & neck cancer in our communities.

KEYWORDS: Cancer Epidemiology, Head & Neck cancer, Incidence, Population.

I. INTRODUCTION

Cancer epidemiology is the study of the distribution and determinants of the likelihood of cancer development. Cancer epidemiology can be used to identify events that increase or decrease cancer incidence in specific populations.

Head and neck cancer is the sixth most common cancer in the world, and the largest burden occurs in developing countries

Head and neck cancer is a group of cancers that starts in the mouth, nose, throat, larynx, sinuses, or salivary glands etc. The annual incidence of head and neck cancers worldwide is more than 550,000 cases with around 300,000 deaths each year. Male to female ratio ranges from 2:1 to 4:1. About 90% of all head and neck cancers are squamous cell carcinomas (HNSCC). HNSCC is the sixth leading cancer by incidence worldwide. Similarly, 57.5% in Asia and largely in India (20 per 100000 population).

Nepal total 9718 new cancer cases were reported by hospital based national cancer registration programme. Whereas, 1882 (19.3%) were head & neck cancer, which was only 1363 (19.2%) out of 8729 cases in 2011 and 1401 (19.4%) in 2012. Similarly, 1741 (19.9%) and 1868 (20.3%) cases of oral cancer were diagnosed for 2013 and 2014.

It shows cancer Head & neck cancer in Nepal was increasing every year. Exact cancer incidence, death and relevant community based details of cancer in Nepal is unknown to address the need of community based epidemiological study was initiated community based cancer research in 15 districts of four state of Nepal. In this study we focus to determine the current status and trend of head & neck cancer in four province of various districts of Nepal .

This was a descriptive epidemiological study with primary and secondary data analysis of new head & neck cancer that are recorded and collected in different data source institution at 15 districts of four state of Nepal science 1st January to 31st December 2013,2014 & 2015. The data analysis was carried out by using SPSS 19.0 This epidemiological study was undertaken at BP Koirala memorial cancer hospital, Bharatpur, Chitwan, Nepal, Which is



| e-ISSN: 2319-8753, p-ISSN: 2320-6710| www.ijirset.com | Impact Factor: 7.512|

|| Volume 9, Issue 9, September 2020 ||

only national cancer institute of the nation, using primary and secondary data from study area in 2013.2014 & 2015. A total of 2950 cases were of registered during 1st January to 31st December 2015. Whereas, only 511 cases were reported for different site of oral head & neck cancer . Among the reported cases of head & neck cancer topography, larynx C 32 (18.4%) was the first leading cancer site followed by brain C71(19.3%) and Other and unspecified parts of tongue C 02 (10.9%) for both sex. Among the female cases larynx C32 (49.2%) was the leading cancer cases followed by brain C71 (47.7%) and other tongue C 02 (27.8%). Similarly, larynx C 32 (20.0%) was the most common cancer among males, followed by brain C 71 (16.4%) and other tongue 39 (12.5%). Head & neck cancer was relatively increased for 2013 to 2014. Whereas, releatively decreased in 2015.

In recent days, the observation of the data of the oral cancer shows its increasing condition. Even in Nepal, the oral cancer is giving to increasing rapidly. According to the medical science, the poor hygiene, alcohol,tobacco use, HPV infection are its major factors. In this regard,. Cancer epidemiology can be used to identify events that increases or decreases cancer incidence in specific populations.

Cancer is a group of disease characterized by uncontrolled, growth, invasion and spread (Metastasis) of abnormal cells. In cancer, normal mechanism of control of growth and cell division are disturbed. Cancer is synonymous with the term "malignant tumor". Cancer one of the common and most important non-communicable disease increasing around the world with marked rise in the least developed parts of the world. Preventing cancer is an important scientific and public health goal of this century. Cancer is becoming a growing public health problem in our country too.

The global cancer burden is estimated to have risen to 18.1 million new cases and 9.6 million deaths in 2018 (GLOBO CAN 2018). Globally, about 1 in 6 deaths is due to cancer. Approximately 70% of deaths from cancer occur in low-and middle-income countries. ii

Asia accounts for 60% of the world population and half the global burden of cancer. The incidence of cancer cases is estimated to increase from 6.1 million in 2008 to 10.6 million in 2030. According to WHO, India has a cancer mortality rate of 79 per 100,000 deaths and accounts for over 6 percent of total deaths. These numbers are very close to those of high-

Table: 1 Area and population covered by this study

S.N.	District's Name	Male	Female	Total
1	Chitwan	298400	297681	658114
2	Makwanpue	243921	238877	528160
3	Bara	361920	347026	708947
4	parsa	322851	305630	628481
5	Nawalparasi	350017	352031	702048
6	Rupandehi	456337	445481	901818
7	Kapilvastu	306323	295986	602309
8	Dhading	204735	207581	412317
9	Gorkha	165830	175941	341771
10	Myagdi	65686	69928	135613
11	Tanahun	183533	195926	378559
12	Baglung	154590	166206	320796
13	Parbat	89095	95287	320796
14	Kaski	235364	240541	475905
15	Mustang	9017	8146	17163
Total		3447619	3442268	6889887

Total population of Nepal: 2,66,208,09(2068) Total Population of Project area: 68,89,887 Coverage25.88% of total population



 \parallel Volume 9, Issue 9, September 2020 \parallel

Table: 2 Distribution of cancer cases by districts for study area

		Distribution of can						
	Districts		Year					
			2013		2014			
		#	%	#	%			
1	Baglung	96	3.9	129	4.9			
2	Bara	165	6.7	176	6.7			
3	Chitawan	359	14.5	424	16.2			
4	Dhading	157	6.4	183	7.0			
5	Gorkha	153	6.2	150	5.7			
6	Kapilbastu	95	3.8	107	4.1			
7	Kaski	323	13.1	325	12.4			
8	Makawanpur	180	7.3	180	6.9			
9	Mustang	10	0.4	15	0.6			
10	Myagdi	44	1.8	47	1.8			
11	Nawalparasi	241	9.8	248	9.5			
12	Parbat	89	3.6	93	3.5			
13	Parsa	123	5.0	146	5.6			
14	Rupandehi	303	12.3	259	9.9			
15	Tanahu	131	5.3	138	5.3			
Total		2469	100.0	2620	100.0			

Table: 3 Distribution of cases by districts and sex of study area for 2015

		Distribution of cas	ses by distri	cts and sex	,		
S.N.	Districts		Male		Female		Total
		#	%	#	%	#	%
1	Baglung	64	5.0	88	5.2	152	5.2
2	Bara	68	5.4	101	6.0	169	5.7
3	Chitwan	176	13.8	198	11.8	374	12.7
4	Dhading	103	8.1	114	6.8	217	7.4
5	Gorkha	94	7.4	126	7.5	220	7.5
6	Kapilvastu	67	5.3	79	4.7	146	4.9
7	Kaski	154	12.1	215	12.8	369	12.5
8	Makwanpur	102	8.0	111	6.6	213	7.2
9	Mustang	6	0.5	12	0.7	18	0.6
10	Myagdi	20	1.6	43	2.6	63	2.1
11	Nawalparasi	104	8.2	172	10.2	276	9.4
12	Parbat	55	4.3	68	4.1	123	4.2



| e-ISSN: 2319-8753, p-ISSN: 2320-6710| <u>www.ijirset.com</u> | Impact Factor: 7.512|

|| Volume 9, Issue 9, September 2020 ||

13	Parsa	69	5.4	82	4.9	151	5.1
14	Rupendehi	117	9.2	165	9.8	282	9.6
15	Tanahun	72	5.7	105	6.3	177	6.0
	Total	1271	100.0	1679	100.0	2950	100.0

II. MATERIALS AND METHODS

This study carried out in the cancer registry unit of B. P. Koirala Memorial Cancer Hospital, Bharatpur, Chitwan, Nepal. The information of all new cancer cases were collected, check, edit and entered in computer using proprietory software. In order to avoid multiple entry of the cases, the data base were verified by name, age, sex, address, topography and morphology of cancer cases. All multiple entries were deleted from the data base. Then, the data was coded according to ICD-03 and ICD-10 iv and proceed for analysis by using SPSS 19.0.

Limitation of the study

Epidemiological Study science 1st January to 31st December 2013, 2014 & 2015 for 15 District of defined geographical region and population.

Data Sources

District Hospital, Medical college and other hospitals.

District Health and Public Health Office and other relevant organizations.

DDC /VDC/Municipalit, i.e. office of vital event registration.

Privet hospitals, Diagnostic lab, hospice etc.

Inclusion Criteria:

The inclusion criteria of cancer cases were set up by National Cancer Registry as follows:

Those Cases living permanently in the study area, as per define population in census.

Diagnosed cases of cancer including mortality in the study year are included.

The minimum diagnostic criteria included tissue diagnosis (HPE or cytology)



|| Volume 9, Issue 9, September 2020 ||

III. RESULT

Table: 4 Head & neck cancer cases for 2013

		Head & neck cancer cases by topography and sex				
S.N.	ICD-	, 1 C 1 7		201	3	
	10- 10	Topography	1	Sex	T.4.1	%
	10		Male	Female	Total	
1	C 32	Larynx	59	35	94	3.8
2	C 71	Brain	33	37	70	2.8
3	C 02	Other & unspecified parts of tongue	30	15	45	1.8
4	C 73	Thyroid	11	26	37	1.4
5	C 06	Other & unspecified parts of mouth	23	8	31	1.2
6	C 12	Pyriform sinus	25	2	27	1.0
7	C 11	Nasopharynx	16	8	24	0.9
8	C 04	Floor of mouth	11	3	14	0.5
9	C 09	Tonsil	9	2	11	0.4
10	C 31	Accessory sinus	6	4	10	0.4
11	C 05	Palate	7	2	9	0.3
12	C 70	Meninges	3	6	9	0.3
13	C 00	Lip	8	0	8	0.3
14	C 30	Nasal cavity & middle ear	3	5	8	0.3
15	C 10	Oropharynx	6	0	6	0.2
16	C 08	Other & unspecified major salivary glands	4	1	5	0.2
17	C 14	Other & ill defined sites in lip oral cavity & mouth	4	1	5	0.2
18	C 69	Eye & adnexa	0	5	5	0.2
19	C 03	Gum	1	2	3	0.1
20	C 01	Base of tongue	2	0	2	0.08
21	C 13	Hypopharynx	1	1	2	0.08
22	C33	Trachea	1	0	1	0.04
23	C 75	Other endocrine glands & related structure	1	0	1	0.04
24	***	Total cases of head & neck cancer	264	163	422	17.0
25	***	Other cancers	849	1193	2047	82.9
		Cancer cases for the year	1113	1356	2469	100.0

Table: 5 Head & neck cancer cases for 2014

		Head & neck cancer cases by topography ar	nd sex for 20	014			
S.N.			2014				
	ICD- 10	Topography		Sex	T. 4.1	%	
	10		Male	Female	- Total		
1	C 32	Larynx	58	27	85	3.2	
2	C 71	Brain	39	34	73	2.7	
3	C 02	Other & unspecified parts of tongue	33	19	52	1.9	
4	C 73	Thyroid	10	37	47	1.7	
5	C 06	Other & unspecified parts of mouth	30	11	41	1.5	
6	C 12	Pyriform sinus	27	1	28	1.0	
7	C 69	Eye & adnexa	14	9	23	0.8	
8	C 30	Nasal cavity & middle ear	9	8	17	0.6	
9	C 11	Nasopharynx	8	8	16	0.6	
10	C 05	Palate	7	5	12	0.4	
11	C 03	Gum	7	3	10	0.3	
12	C 00	Lip	4	4	8	0.3	



|| Volume 9, Issue 9, September 2020 ||

13	C 04	Floor of mouth	4	4	8	0.3
14	C 10	Oropharynx	6	2	8	0.3
15	C 31	Accessory sinus	4	4	8	0.3
16	C 08	Other & unspecified major salivary glands	5	1	6	0.2
17	C 09	Tonsil	2	2	4	0.1
18	C 70	Meninges	0	4	4	0.1
19	C 13	Hypopharynx	3	0	3	0.1
20	C 14	Other & ill defined sites in lip, oral cavity & mouth	2	0	2	0.07
21	C 75	Other endocrine glands & related structures	0	2	2	0.07
22	C 01	Base of tongue	1	0	1	0.03
23	***	Total cases of head & neck cancer	273	185	458	17.4
24	***	Other cancer	815	1347	2162	82.5
		Total cancer cases for the year	1088	1532	2620	100.0

Table: 6 Head and neck cancer for 2015

		Distribution of head and neck cancer cases b	y topogra	phy and sex	ī	
S.N.	ICD -	Topography		Sex	,	Total
	10		Male	Female	#	%
1	C 00	Lip	4	0	4	0.1
2	C 02	Other and unspecified parts of tongue	39	17	56	1.9
3	C 03	Gum	11	3	14	0.5
4	C 04	Floor of mouth	2	1	3	0.1
5	C 05	Palate	7	0	7	0.2
6	C 06	Other and unspecified parts of mouth	28	8	36	1.2
7	C 07	Parotid gland	7	5	12	0.4
8	C 08	Other and unspecified major salivary glands	2	3	5	0.2
9	C 09	Tonsil	8	3	11	0.4
10	C 10	Oropharynx	2	2	4	0.1
11	C 11	Nasopharynx	14	6	20	0.7
12	C 12	Pyriform sinus	29	8	37	1.3
13	C 13	Hypopharynx	2	2	4	0.1
14	C 14	Other and ill-defined sites in lip, oral cavity and pharynx	4	0	4	0.1
15	C 30	Nasal cavity and middle ear	10	7	17	0.6
16	C 31	Accessory sinus	4	4	8	0.3
17	C 32	Larynx	62	37	99	3.4
18	C 69	Eye and adnexa	7	1	8	0.3
19	C 70	Meninges	1	3	4	0.1
20	C 71	Brain	51	45	96	3.3
21	C 73	Thyroid gland	9	38	47	1.6
22	C 75	Other endocrine glands and related structures	1	0	1	0.0



| e-ISSN: 2319-8753, p-ISSN: 2320-6710| <u>www.ijirset.com</u> | Impact Factor: 7.512|

|| Volume 9, Issue 9, September 2020 ||

23	C 76	Other and ill-defined sites of head, face & neck	6	8	14	0.4
24	***	Total cases of head & neck cancer	310	201	511	17.32
25	***	Other cancer cases	961	1478	2439	82.6
		Total cancer cases for the year	1271	1679	2950	100.0

IV. DISCUSSION

This epidemiological study was undertaken at BP Koirala memorial cancer hospital, Bharatpur, Chitwan, Nepal, Which is only national cancer institute of the nation, using primary and secondary data from study area in 2013.2014 & 2015. A total of 2950 cases were of registered during 1st January to 31st December 2015. Whereas, only 511 cases were reported for different site of oral head & neck cancer . Among the reported cases of head & neck cancer topography, larynx C 32 (18.4%) was the first leading cancer site followed by brain C71(19.3%) and Other and unspecified parts of tongue C 02 (10.9%) for both sex. Among the female cases larynx C32 (49.2%) was the leading cancer cases followed by brain C71 (47.7%) and other tongue C 02 (27.8%). Similarly, larynx C 32 (20.0%) was the most common cancer among males, followed by brain C 71 (16.4%) and other tongue 39 (12.5%). Head & neck cancer was relatively increased for 2013 to 2014. Whereas, releatively decreased in 2015 due to the massive earthquake.

V. CONCLUSION

Among the reported cases of head & neck cancer, larynx C 32 (18.4%) was the first leading cancer site followed by brain C71(19.3%) and Other and unspecified parts of tongue C 02 (10.9%) for both sex.

REFERENCES

1.International Classification of Disease for oncology, WHO/IARC 3rd Edition, Lyon France, 2000
2.National Cancer Registry Programme, Report of Hospital Based National Cancer Registry 2003, B.P. Koirala
Memorial Cancer Hospital Bharatpur, Chitwan, Nepal
3. National Cancer Registry Programme, Report of Hospital Based National Cancer Registry 2004, B.P. Koirala
Memorial Cancer Hospital Bharatpur, Chitwan, Nepal
4. National Cancer Registry Programme, Report of Hospital Based National Cancer Registry 2005, B.P. Koirala
Memorial Cancer Hospital Bharatpur, Chitwan, Nepal
5.National Cancer Registry Programme, Report of Hospital Based National Cancer Registry 2006, B.P. Koirala
Memorial Cancer Hospital Bharatpur, Chitwan, Nepal
6.National Cancer Registry Programme, Report of Hospital Based National Cancer Registry 2007, B.P. Koirala
Memorial Cancer Hospital Bharatpur, Chitwan, Nepal
7. National Cancer Registry Programme, Report of Hospital Based National Cancer Registry 2008, B.P. Koirala
Memorial Cancer Hospital Bharatpur, Chitwan, Nepal
8. National Cancer Registry Programme, Report of Hospital Based National Cancer Registry 2009, B.P. Koirala
Memorial Cancer Hospital Bharatpur, Chitwan, Nepal
9.National Cancer Registry Programme, Report of Hospital Based National Cancer Registry 2010, B.P. Koirala
Memorial Cancer Hospital Bharatpur, Chitwan, Nepal
10. National Cancer Registry Programme, Report of Hospital Based National Cancer Registry 2011, B.P. Koirala
Memorial Cancer Hospital Bharatpur, Chitwan, Nepal
11. National Cancer Registry Programme, Report of Hospital Based National Cancer Registry 2012, B.P. Koirala
Memorial Cancer Hospital Bharatpur, Chitwan, Nepal
12. National Cancer Registry Programme, Report of Hospital Based National Cancer Registry 2013, B.P. Koirala
Memorial Cancer Hospital Bharatpur, Chitwan, Nepal
13. National Cancer Registry Programme, Report of Hospital Based National Cancer Registry 2014, B.P. Koirala
Memorial Cancer Hospital Bharatpur, Chitwan, Nepal
14. National Cancer Registry Programme, Report of Hospital Based National Cancer Registry 2015, B.P. Koirala
Memorial Cancer Hospital Bharatpur, Chitwan, Nepal
15. Global cancer statistics 2018



| e-ISSN: 2319-8753, p-ISSN: 2320-6710| www.ijirset.com | Impact Factor: 7.512|

|| Volume 9, Issue 9, September 2020 ||

16.Edge, Stephen B. (2010). AJCC cancer staging manual. American Joint Committee on Cancer (7th ed.). New York: Springer. ISBN 9780387884400. OCLC 316431417.

- 17. E., Marx, Robert (2003). Oral and maxillofacial pathology : a rationale for diagnosis and treatment. Stern, Diane. Chicago: Quintessence Pub. Co. ISBN 978-0867153903. OCLC 49566229.
- 18. "Head and Neck Cancers". CDC. 2019-01-17. Retrieved 2019-03-10.
- 19. Cancer today". gco.iarc.fr. Retrieved 9 June 2019.
- 20. "USCS Data Visualizations". gis.cdc.gov. Retrieved 2019-03-10.









Impact Factor:

INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY







📵 9940 572 462 🔊 6381 907 438 🖂 ijirset@gmail.com

