



e-ISSN: 2319-8753 | p-ISSN: 2347-6710

IJIRSET

International Journal of Innovative Research in
SCIENCE | ENGINEERING | TECHNOLOGY

INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 12, Issue 10, October 2023

ISSN INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA

Impact Factor: 8.423

9940 572 462

6381 907 438

ijirset@gmail.com

www.ijirset.com

A Review of the Structure of Water

Md. Shamsuddin

Maharana Pratap Govt. P.G. College, Chittorgarh (Raj.), India

ABSTRACT: Water is a remarkable component with various unique features. It is being argued that utilizing the "anomalous" characteristics of water could result in intriguing possibilities in a variety of fields, including medical science, engineering, and biochemistry. The powerful bonding of hydrogen found in water only partially explains the beginnings of these odd features. Furthermore, the structure and dynamics of water are still insufficiently understood, and novel hypotheses have emerged and been hotly contested in the modern era. Water's qualities as a solvent are the main topic of this account. Boasting dissolution spanning more than 10 powers of severity, water is able to dissipate a wide variety of compounds. The basic distinction between polar, or hydrophilic, and apolar, or hydrophobic, solutes is defined by variations in solubility. The structure, equilibrium, and functionality of biological macromolecules are significantly influenced by this crucial differential. The bidirectional rotational arrangement of the H-O, OH-bond angles determines the magnitude of the bond between hydrogen molecules.

KEYWORDS: Water molecules, structure, hydrophobic, Hydrogen bonds, Solvent

I. INTRODUCTION

Water is a special, pervasive element that is essential to all creatures that exist. Since ancient times, intellectuals, ecologists, and geologists have been fascinated by its origins and qualities. Despite years of intensive research, scientists are still interested in water because of how little is known about it. This is mainly because water has various chemical as well as physical characteristics that are anomalous. Certain of the water's special qualities are essentially necessary for living, whereas others have a significant impact on the form and dimension of living things, the way they function, and the physical restrictions they must adhere to. ^[1] Water additionally has a major influence on the chemical composition and luminescence of the atmosphere. Numerous members of the atmosphere's water family, which ranges from individual molecules of hydrogen and oxygen through a concentrated section, i.e. fluid, water, and ice, are descended from the water particle. The consequences of atmospheric water disintegration and consolidation are particularly significant for biological life on Terra. Short-wave UV radiation (as well as those lower than 242 nm) can cause the hydrogen atom to separate from the water molecule. ^[2]

Although oxygen Herzberg attenuation severely reduces UV light at lower elevations; direct transpiration of water only occurs efficiently in the higher stratosphere. Because of this, molecular H₂O changes and breakdowns are common under 50 km. Water collects mostly in the upper troposphere, where vaporized water's relative density is substantial. The accumulation results from comparatively weak intermolecular connections among water-water and water-foreign molecules of gas that form in close proximity. One of the primary aspects that defines how water molecules associate is the so-called bond between hydrogen molecules^[3]. It's noteworthy to note that the formation of hydrogen bonds is essential for all forms of organic life since they allow for the complex nucleotic bases and acids' distinctive reorganizations. Since then, additional information has been gained, largely as a result of spectroscopy and thermodynamic research, concerning the molecular composition and characteristics of water.

Machine modeling, an additional recent discipline, has also contributed since it has advanced to the point of research in water where it may be employed to interpret observations and replicate qualities that are not easily accessible by trial. Today, many of the fundamental physical characteristics of water can be at least partially described by means of their molecular and structural configuration.

The Molecular Structure Of Water

The two moderately positively charged hydrogen atoms are joined by an extremely electronegative oxygen particle to form water. As a consequence, an extremely polar molecule is created, one that can contribute H-bonds via strongly charged hydrogen atoms and acquire H-bonds from inversely charged O atoms, among other powerful electrostatic forces. Finding the framework for Ice 1h was an important achievement from an architectural standpoint.

The common form of ice, known as ice 1h, can be observed at intermediate pressure as well as temperature levels. The now-famous tetrahedral ice lattice framework was discovered by Bragg. At an angle of 2.75, every water molecular structure has four hydrogen-bound first neighbors, two of which act as H-bond donors and two of which act as H-bond acceptors.^[4]

The four companions are positioned with 4.5 units of space between them at the edges of a tetrahedron. Since liquid water's organization has not yet been properly defined, our understanding of ice 1h, a relatively open framework with an elevated level of axial and circular ordering, has been greatly influenced. Since each molecular structure contains two hydrogen atoms, it makes sense that each molecule in fluid water ideally ought to contribute two H-bonds, just like in ice.^[5] It is less evident why accepting two H-bonds is the best option. In fact, a lot of research continues to center on the quantity and structure of H-bonds produced by liquid water.

Structure of Water:

Water is regarded as one of the most versatile solvents for large number of electrolytes, and it is regarded as amphiprotic, dipolar, protic or neutral amphiprotic solvent having the physical and molecular property as given in table I

Property of water	Value
Melting point	273.15K
Boiling point	372.15K
Critical temperature	647.55K
Dielectric constant (at 298 K)	78.54K
Viscosity Coefficient (at 298 K)	0.900 cp.
Dipole movement	1.8834 D
DN	18 (33)
AN	54.8

The identification of O-O, H-H, and O-H radial dispersion coefficients ($g(r)$) by X-ray and neutron dispersion investigations was a breakthrough in our comprehension of the molecular makeup of water as a fluid.²⁻⁴ Surprisingly, data demonstrate that a significant portion of the tetrahedral ice lattice survives in fluid water: according to the oxygen-oxygen zonal dispersion (local concentration) measure $g_{OO}(r)$, the closest neighbors have a density crest at 2.8-2.9Å, which is just marginally further away than in ice 1h. Additionally, there exists a remnant intensity spike at 4.5-4.7 that is quite wide but still indicative of the initial collaboration shell's tetrahedral arrangement of the fluids. Nevertheless, the axial dispersion characteristics show that there are significant variations across liquid and ice.^[6]

Essentially derivable using X-ray/neutron scattering tests, circumferential dispersion functions have produced an abundance of knowledge for a variety of temperatures as well as pressures. Fewer are known regarding the molecular alterations that occur with solvation as a result of these investigations, which use neutron dispersion and hydrogen/deuterium isotope replacement to conduct accurate experimental research.^[7,8] For instance, it seems that the separation among the closest neighboring fluids surrounding ionic and apolar solvents is fairly consistent surrounding 2.75-2.9.

II. DISCUSSION

Water is used as a versatile solvent due to the hydrogen bonding effect. Water molecules contain hydrogen atoms and two free lone pairs of electrons, which enable to form four hydrogen bonds that are tetrahedrally arranged. Due to this position of the free electron pairs and two hydrogen atoms.

There has been great controversy regarding the structure of water, although there has been major breakthrough in regrading structure of water Yet no definite structure of water molecules has been provided. The structure of water has been given by the collective venture of many workers. Many experimental advances and models have been proposed for the structure of water molecules. On the basis of X-ray studies, Bernal and Fowler^[9] have calculated that ice is tetrahedrally coordinated, having an open structure, a short range, and a short time. Liquid water retains the tetrahedral structure of ice: Morgan and warren^[10] found that the coordination number of water at 277.5 K is 4.4 by diffraction studies, but at a little distance away, 1.63 times the O-O distances in ice, the coordination number is 4, and hydrogen bonds are tetrahedrally arranged. The immediate environment of a water molecule is liquid water consisting of the nearest and the second nearest, which is very ice like, but this environment persists over a distance of two times the O-O distances, so from the above discussion, water has an open structure and is full of cavities.

'Interstitial' model has been proposed for water by Samoilov^[11] at a lower temperature, cavities are filled by monomeric, non-hydrogen bonded water molecules. Later on, Mikhailov^[12] has been expressed a similar view. Pauling^[13] on the other hand, has considered liquid water as something resembling the clathrate compounds analogous to the gas hydrates formed between methane and water. Water forms a hydrogen bonded cage around monomeric water molecules.

NEMETHY and Scheraga^[14] however, considered water to consist of a two-state mixture. Here, liquid water is regarded as a mixture of open, empty, lighter clusters and a denser fluid composed of hydrogen bonded molecules. In addition to hydrogen bonding, they also considered that there is some other kind of strong water-water interaction that has not yet been cleared. Evidence for such a structural representation of liquid water was furnished later on by the work of Walrafen^[15]. A modification of this two state model for liquid water, the so called 'flickering cluster model' was proposed by Frank and Coworkers^[16,17], who thought that hydrogen bonding in water is an essentially cooperative process very well suited to form clusters. Quantitatively, the flickering cluster model involves a large number of water molecules that are momentarily all hydrogen-bonded with one another; with one to four bonds per molecule. There is also possibility of a few non-hydrogen bonded water molecules are also present. A water molecule participating in the formation of one bond is likely to participate in the formation of another bond also, since the bond formation is cooperative. The opposite process is true as well. If, as a result of thermal agitation one bond is broken, several more such bonds are also broken, resulting in the cluster 'flicker', the cluster being of several shapes and sizes and have half is the order of 10^{-10} to 10^{-11} sec.

Scherer and Coworkers^[18] on the other hand, have proposed a mixture model for liquid water that may be regarded as a mixture of a small number of distinct molecular species having a variable number of hydrogen bonds per molecule. In another two-state mixture model of water, Marchi and Eyring^[19] have proposed that these are two distinctively separate species existing in liquid water. These are the freely rotating monomeric water molecules and the hydrogen bonded ice like water. This model has been invoked to explain the maximum density of liquid water at 277.15K.

Scheraga and coworkers^[20] and Stillinger and coworkers^[21,22] have given theories regarding slow formation of tetrahedral hydrogen bonds between neighboring water molecules. But the hydrogen bonds are sometimes bent away from the ideal tetrahedral hydrogens bonds are sometimes bent away from the ideal tetrahedral direction and large clusters of anomalous density is not formed.

In water, the three hindered rotatory motions change significantly with temperature and pressure. Thus affecting the specific heat. But the exact relationship between specific heat and rotatory motion is not understood since the macrodynamic structure of water seems to be complex^[23].

In addition to the above model of water, many newer models regarding the geometrical structure of water have been proposed^[24-29]. Therefore, a large number of models for liquid water have been proposed, but in many respects, the proposed structures are contradictory to one another [28].

Based upon the above explanation, the probable structure of water may be taken as a 'tetrahedral' quasi-crystalline structure constantly changing and giving a disordered structure^[29] with the production of monomeric water molecules as well.

Water is supposed to behave as an associated liquid at its normal point because its Trouton's constant is higher than 21 CalK^{-1} . The donor number of water is not surprising high; it is 18, but the acceptor number is 54.8, which is a very high value.

Based upon these features of water, it has outstanding capabilities for solvent anion, which is an electron donor. It also has the capability of solvent cation, but it is small in comparison to solvation of anion. Several water molecules are able to be associated with cation, and several water molecules are able to be associated with a cation. A cooperative process starts with a much higher donor number, about 33, rather than 18. The high dielectric constant is also due to the cooperative alignment of the dipole, which partly causes high hydrogen bonding, or it may be said that hydrogen bonding is one of the best features present in water molecules. Due to this, it is regarded as the most versatile solvent. The high surface tension of water is also dependent on this cause.

REFERENCES

- Chlek P. and Geldart D.J.W. Water vapor dimers and atmospheric absorption of electromagnetic radiation. *Geophys. Res. Letters*. 1997; 24(16) 2015-2018.
- Deepak A., Wilkerson T.D., and Ruhnke L.H., Eds. *Atmospheric Water Vapor*. 1980; 695 pp. New York: Academic Press.
- Eisenberg D. and Kauzmann W., *The Structure and Properties of Water*, 1969, 296 pp. Oxford: Clarendon Press.
- Schirò G, Weik M. Role of hydration water in the onset of protein structural dynamics. *J. Phys. Condens. Matter*. 2019;31(46):463002.
- Rahman A. and Stillinger F., *Molecular Dynamics Study of Liquid Water* *Journal of Chemical Physics*. 1971;55: 3336-3359.
- Henderson, L.J., *The Fitness of the Environment: An Inquiry into the Biological Significance of the Properties of Matter*. 1913; New York: Macmillan
- Xu Y, Havenith M. Perspective: watching low-frequency vibrations of water in biomolecular recognition by THz spectroscopy *J. Chem. Phys.* 2015;143(17):170901.
- Turner J., Soper A., and Finney J. Ionic versus apolar behavior of the tetramethylammonium ion in water. *J. Chem. Phys* 1995;102:5438-5443.
- Bernal J. D. and Fowler R. H. 1933 *J. Chem. Phys.* 1 515.
- Morgan J., Warren BE. X-ray analysis of the structure of water. *The Journal of Chemical Physics*. 1938 Nov 1;6(11):666-73.
- Samoilov OY. The coordination number in the structure of certain liquids *Zh. Fiz. Khim.*. 1946;20:1411.
- Mikhailov VA, Ponomareva LI. Structure and thermodynamic properties of aqueous solutions of nonelectrolytes: I. Description of the model. *Journal of Structural Chemistry*. 1968 Jan;9(1):8-15.
- Pauling, L., *The Nature of the Chemical Bond—An Introduction to Modern Structural Chemistry*. 3rd Edition, Cornell University Press, Ithaca, New York. 1960:10-13.
- SCHERAGA HA, NEMETHY G, STEINBERG IZ. The contribution of hydrophobic bonds to the thermal stability of protein conformations *J Biol Chem*. 1962 Aug;237:2506-8. PMID: 14498257.
- Walrafen, George E., "Raman Spectral Studies of Water Structure." *Journal of Chemical Physics* 40 (1964): 3249-3256.
- Frank, Henry S., and Wen Yang Wen. "Ion-solvent interaction. Structural aspects of ion-solvent interaction in aqueous solutions: a suggested picture of water structure" *Discussions of the Faraday Society* 24 (1957): 133-140.
- Scheraga HA. Tyrosyl-carboxylate ion hydrogen bonding in ribonuclease *Biochemistry Biophysics Acta*. 1957;23:196-197.
- Liang K, Eyring H, and Marchi RP SIGNIFICANT STRUCTURE THEORY OF BINARY LIQUID MIXTURES: CARBON TETRACHLORIDE AND CYCLOHEXANE *Proc Natl Acad Sci U S A*. 1964 Oct;52(4):1107-14. doi: 10.1073/pnas.52.4.1107. PMID: 16591227; PMCID: PMC300402.
- Hagler AT, Scheraga HA, and Nemethy G. Structure of liquid water. *Statistical thermodynamic theory* *The Journal of Physical Chemistry*. 1972;76: 3229-3243. DOI: 10.1021/J100666A022.
- Rahman, Aneesur, and Frank H. Stillinger "Molecular Dynamics Study of Liquid Water." *Journal of Chemical Physics* 55 (1971): 3336-3359.
- Kell, G.S. *Thermodynamic and Transport Properties of Fluid Water*. In: Franks, F. (eds.) *The Physics and Physical Chemistry of Water*. Water, 1972, vol. 1. Springer, Boston, MA. https://doi.org/10.1007/978-1-4684-8334-5_10.
- Frank HS, Wen WY, and PITTSBURGH UNIV PA Ion-solvent Interaction III Structural Aspects of Ion-solvent Interaction in Aqueous Solutions: A Suggested Picture of Water Structure 1957.
- H. Eugene Stanley, J. Teixeira; Interpretation of the unusual behavior of H₂O and D₂O at low temperatures: Tests of a percolation model. *J. Chem. Phys.* 1 October 1980; 73 (7): 3404-3422. <https://doi.org/10.1063/1.440538>
- Stanley, H. E., Blumberg, R. L., and Geiger, A. Gelation models of hydrogen bond networks in liquid water, *Phys. Rev. B*. 1981;28, 1626-1629.
- Behlen, Fred M., Rice, S. A. Intersystem crossing in cold isolated molecules of naphthalene *The Journal of Chemical Physics*. 1981;75 (12). 5672- 5684 doi:10.1063/1.442004.
- Meijer, P.H.E., Kikuchi, Ryoichi, and Royen, Eddy Extended Lattice Model of Water; Two-Sublattice, Two-Orientation Model *Physica A-Statistical Mechanics and Its Applications - PHYSICA A*. 1982;115. 124-142. 10.1016/0378-4371(82)90131-5.
- J. V. Levenekkers, *J. Chem. Soc., Faraday Trans.* 1988;2, 84, No. 2, 397-411.
- J. Lee Kavanau, *Water and Solute-Water Interactions*, 1964.
- Keith James Laidler, *Chemical Kinetics*, Second Edition, 1965.



INNO SPACE
SJIF Scientific Journal Impact Factor
Impact Factor: 8.423



ISSN INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

9940 572 462 6381 907 438 ijirset@gmail.com



www.ijirset.com

Scan to save the contact details