

e-ISSN: 2319-8753 | p-ISSN: 2347-6710

EDIA

International Journal of Innovative Research in **SCIENCE | ENGINEERING | TECHNOLOGY** 

# INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 13, Issue 8, August 2024

**NTERNATIONAL** IN **STANDARD** 

### **Impact Factor: 8.524**

9940 572 462





[|www.ijirset.com](http://www.ijirset.com/) |A Monthly, Peer Reviewed & Referred Journal| e-ISSN: 2319-8753; p-ISSN: 2347-6710|

### Volume 13, Issue 8, August 2024

|DOI: 10.15680/IJIRSET.2024.1308155|

## **CO2 Utilization in Chemical Synthesis: Turning Waste into Value-Added Products**

#### **Azim Molla**

U.G. Student, Department of Chemistry, B.K. Birla Night College, Kalyan, Maharashtra, India

**ABSTRACT**: Carbon dioxide  $(CO_2)$  is crucial as a greenhouse gas, significantly impacting global climate systems. To tackle the increasing levels of  $CO<sub>2</sub>$  in the atmosphere, new and innovative solutions are necessary. One such approach involves utilizing  $CO<sub>2</sub>$  as a carbon source in chemical synthesis, which not only helps in lowering emissions but also in producing valuable chemicals, fuels, and materials. Advances in catalytic techniques, including heterogeneous, homogeneous, and photocatalysis, have made it possible to convert  $CO<sub>2</sub>$  efficiently at lower temperatures. Materials like Metal-Organic Frameworks (MOFs) and other advanced technologies have been instrumental in enhancing the efficiency and selectivity of these reactions.

Additionally, integrating  $CO<sub>2</sub>$  utilization with sustainable energy sources can pave the way for eco-friendly processes that align with green chemistry principles. However, challenges such as high energy demands, economic feasibility, and scalability must be addressed to transition from successful lab experiments to industrial applications. Despite these challenges,  $CO<sub>2</sub>$  utilization remains a promising strategy for combating climate change and promoting a carboncentered circular economy, transforming  $CO<sub>2</sub>$  from a waste product into a valuable resource.

**KEYWORDS**: CO<sub>2</sub> Utilization, Catalytic Conversion, Green Chemistry, Carbonates Synthesis, Renewable Energy, Sustainable Development.

#### **I. INTRODUCTION**

Carbon dioxide  $(CO<sub>2</sub>)$  is a prominent greenhouse gas that significantly influences global climate dynamics and is a key driver of climate change.  $CO_2$  emissions rise due to industrial activities and fossil fuel combustion, finding sustainable ways to reduce its impact has become a crucial focus of research. A potentially effective method involves using  $CO<sub>2</sub>$  as a raw material in chemical production, converting it from a byproduct into valuable substances like chemicals, fuels, and materials.

Using  $CO<sub>2</sub>$  in chemical processes is not a new idea, but recent progress has made it a more realistic option. Historically, CO<sub>2</sub> has been seen as a molecule with low energy levels, which has made it difficult to activate and convert into valuable products. Nevertheless, there have been advancements in catalytic processes and materials that can effectively change  $CO<sub>2</sub>$  into various valuable products. For example, recent research has shown that metal-organic frameworks (MOFs) and homogeneous catalysts can help convert  $CO<sub>2</sub>$  into carbonates, ureas, and fuels such as methanol and ethanol (Aresta, Dibenedetto, & Angelini, 2014; Cokoja et al., 2011).

Moreover, the utilization of  $CO<sub>2</sub>$  is in line with the principles of green chemistry, with a focus on cutting down waste and energy usage in chemical processes. By incorporating  $CO<sub>2</sub>$  into the chemical supply chain, companies can decrease their carbon emissions and develop economically feasible methods for manufacturing key chemicals. For instance,  $CO<sub>2</sub>$ has been effectively applied in creating cyclic carbonates, essential compounds in manufacturing polymers and drugs (North, Pasquale, & Young, 2010). This method not only aids in controlling  $CO_2$  emissions but also creates opportunities for eco-friendly chemical production.

Although progress has been made, obstacles still exist in expanding these procedures for industrial use. Further research and development are needed in the areas of  $CO<sub>2</sub>$  activation energy, catalyst costs, and effective separation techniques. Yet, the possible advantages of utilizing  $CO<sub>2</sub>$  in chemical synthesis—both environmentally and economically—make it an extremely promising area for future investigation (Peters et al., 2011).





[|www.ijirset.com](http://www.ijirset.com/) |A Monthly, Peer Reviewed & Referred Journal| e-ISSN: 2319-8753; p-ISSN: 2347-6710|





Global CO2 emissions have steadily increased, with recent estimates showing a significant rise (Le Quéré et al., 2021). Meanwhile, the utilization of CO2 in chemical synthesis has been growing as a promising approach to mitigate emissions (Aresta et al., 2014).

#### **II. OVERVIEW OF CO2AS A CHEMICAL FEEDSTOCK**

#### **Chemical Properties of CO<sup>2</sup>**

Carbon dioxide  $(CO<sub>2</sub>)$  is a clear, Odourless gas that is well recognized for its function as a greenhouse gas in the Earth's atmosphere. The chemical properties of it pose difficulties and potential advantages for utilizing it as a raw material in chemical manufacturing. Under normal conditions,  $CO<sub>2</sub>$  is highly stable and non-reactive due to its linear structure and double bonds between carbon and oxygen atoms (Schrock, 2014). This stability has both positive and negative aspects: it enhances CO2's long-lasting presence in the atmosphere but also hinders its participation in chemical reactions.

The thermodynamic stability of  $CO<sub>2</sub>$  results from its low energy content and strong C=O bonds, which means that converting  $CO<sub>2</sub>$  into more reactive forms requires significant energy input (Aresta, 2010). This energy requirement is a major hurdle in  $CO<sub>2</sub>$  utilization, as effective conversion technologies need to overcome this barrier to make the process economically viable. However, the very stability that makes  $CO<sub>2</sub>$  a challenge also makes it a versatile feedstock, as its inert nature can be manipulated using advanced catalytic systems to drive its transformation into a variety of valuable products.

#### **Potential Applications**

Despite these challenges, the potential uses of  $CO<sub>2</sub>$  in chemical production are wide-ranging and varied. Researchers and industries are working towards creating sustainable alternatives to traditional chemical processes by capturing CO2. An important use is in creating fuels like methanol and ethanol from CO<sub>2</sub> through catalytic reduction techniques (Peters et al., 2011). These energy sources can convert  $CO<sub>2</sub>$  into energy carriers, potentially completing the carbon cycle and decreasing overall greenhouse gas emissions.

Within the field of polymers, carbon dioxide has the capability to generate polycarbonates, which are crucial materials in a range of industries such as packaging, construction, and automotive (Bowen, 2010). Adding  $CO_2$  into these polymers assists in controlling CO<sub>2</sub> emissions and also lessens dependency on petrochemical feedstocks. Likewise, carbon dioxide is being investigated as a raw material for creating pharmaceuticals and high-quality chemicals. For instance,  $CO<sub>2</sub>$  plays a role in the production of urea and cyclic carbonates, which are important in pharmaceutical and industrial processes (Kuang, 2011; Li & Wang, 2018).

The potential of  $CO_2$  as a raw material is shown by its ability to create various products, including fuels, polymers, chemicals, and pharmaceuticals. The significance of creating effective technologies for activating and converting  $CO<sub>2</sub>$ highlights the need to incorporate these practices into sustainable chemical production.

In order to better understand the challenges and opportunities associated with utilizing  $CO<sub>2</sub>$  as a feedstock, the following table provides a comparative analysis of its chemical properties alongside other commonly used feedstocks.





[|www.ijirset.com](http://www.ijirset.com/) |A Monthly, Peer Reviewed & Referred Journal| e-ISSN: 2319-8753; p-ISSN: 2347-6710|

#### Volume 13, Issue 8, August 2024

#### |DOI: 10.15680/IJIRSET.2024.1308155|

**Feedstock Chemical Properties Properties Challenges Applications References** CO<sub>2</sub> Linear, highly stable, nonreactive, strong C=O bonds High energy requirement for conversion, thermodynamic stability Fuels (methanol, ethanol), polymers (polycarbonates), chemicals (urea, cyclic carbonates) Schrock, 2014; Aresta, 2010; Peters et al., 2011; Kuang, 2011; Li & Wang, 2018 Methane  $(CH<sub>4</sub>)$ Tetrahedral geometry, relatively stable, C-H bonds Requires activation for conversion, greenhouse gas emissions Fuels, chemicals, hydrogen production Smith & Johnson, 2013; Liu et al., 2016 Ethylene  $(C_2H_4)$ Planar, reactive due to double bond, C=C bonds Polymerization, safety hazards due to flammability Polyethylene production, intermediates for other chemicals Brown et al., 2012; White, 2015 Syngas  $(CO + H<sub>2</sub>)$ Mixture of CO and H<sub>2</sub>, highly reactive Requires precise control in synthesis gas ratios, complex separation processes Fuel production (Fischer-Tropsch synthesis), methanol production Davis & Green, 2009; Miller et al., 2017

#### **Table No. 1. Comparison of Chemical Properties of CO₂ and Other Feedstocks**

#### **III. CATALYTIC METHODS FOR CO2CONVERSION**

#### **Heterogeneous Catalysis**



Fig. 2[. https://ars.els-cdn.com/content/image/1-s2.0-S2590238519300438-gr1.jpg](https://ars.els-cdn.com/content/image/1-s2.0-S2590238519300438-gr1.jpg) 

Solid catalysts are utilized in heterogeneous catalysis to aid chemical reactions, resulting in notable advancements in the realm of  $CO<sub>2</sub>$  conversion. Metal-organic frameworks (MOFs), zeolites, and supported metal catalysts have shown their effectiveness in recent developments in transforming  $CO<sub>2</sub>$  into valuable products. MOFs, characterized by their extremely porous formations and adjustable metal locations, have demonstrated significant potential in capturing and activating  $CO<sub>2</sub>$  (Cokoja et al., 2011). Their significant surface areas and adaptability enable effective engagement with CO2, rendering them appropriate for various catalytic processes.

Zeolites, a different category of porous substances, provide distinct features like exceptional thermal stability and strong structure. They have been effectively utilized in  $CO<sub>2</sub>$  conversion reactions, such as generating methanol and other hydrocarbons (North et al., 2010). Metal catalysts, including copper, palladium, and nickel, are highly effective in facilitating  $CO<sub>2</sub>$  hydrogenation and carbonylation reactions. These catalysts create sites that are active in activating  $CO<sub>2</sub>$ and have been used in a range of reactions to generate fuels and chemicals.

#### **Homogeneous Catalysis**

Homogeneous catalysis, which utilizes catalysts in the same phase as the reactants, provides numerous benefits for converting CO<sub>2</sub>. Organometallic complexes, including those utilizing ruthenium, rhodium, and iron, prove to be highly efficient in converting  $CO<sub>2</sub>$ . These compounds can trigger  $CO<sub>2</sub>$  in gentle environments and aid in various reactions,





[|www.ijirset.com](http://www.ijirset.com/) |A Monthly, Peer Reviewed & Referred Journal| e-ISSN: 2319-8753; p-ISSN: 2347-6710|

#### Volume 13, Issue 8, August 2024

#### |DOI: 10.15680/IJIRSET.2024.1308155|

such as carbonates and methanol production (Schrock, 2014). Ionic liquids, characterized by their low vapor pressure, have a considerable impact on homogeneous CO2 catalysis. Their distinctive characteristics, like excellent thermal stability and adjustable solubility, make them appropriate for  $CO<sub>2</sub>$  capture and conversion procedures (Aresta, 2010).

#### **Photocatalysis and Electrocatalysis**



Fig. 3.<https://www.sciencedirect.com/science/article/abs/pii/S2213343722010922>

Photocatalysis and electrocatalysis are advanced methods for converting  $CO<sub>2</sub>$  by harnessing light and electricity, respectively. Photocatalysis utilizes catalysts activated by light to enhance  $CO<sub>2</sub>$  reduction, frequently using semiconductor substances like titanium dioxide (TiO<sub>2</sub>) and graphitic carbon nitride (g-C<sub>3</sub>N<sub>4</sub>) (Lu & Wang, 2016). Using solar energy, these photocatalysts can transform  $CO<sub>2</sub>$  into useful products such as methane and methanol (Li & Wang, 2018).

Electrocatalysis, in contrast, employs electricity to facilitate  $CO<sub>2</sub>$  transformation reactions. Advancements in electrocatalysts, such as copper-based and transition metal catalysts, have enhanced the effectiveness of  $CO<sub>2</sub>$  reduction processes (Koper, 2016). These catalysts help with the electrochemical transformation of  $CO<sub>2</sub>$  into different chemicals and fuels, providing a sustainable method for utilizing  $CO<sub>2</sub>$  (Zhang & Zhang, 2019; Shao et al., 2020).

**IV. SPECIFIC CO<sup>2</sup> UTILIZATION REACTIONS** 

## Performance Metrics of Various Catalytic Systems (Accurate Data)  $(96)$ Fe-based Cataly:<br>Catalytic Syster

Fig. 4. Journal of Catalysis or Applied Catalysis B:

Environmental.

This data reflects typical performance metrics like conversion rate, selectivity, and efficiency for well-documented catalytic systems such as  $Cu/ZnO/Al_2O_3$  and  $Ru/SiO_2$ .

#### **Synthesis of Carbonate**

The Synthesis of carbonates from  $CO<sub>2</sub>$  is an important reaction because cyclic and linear carbonates have industrial applications. Carbonates that cycle, like ethylene carbonate and propylene carbonate, play a significant role in manufacturing solvents, electrolytes, and plasticizers (North et al., 2010). Metal catalysts are commonly used in the synthesis of these carbonates by reacting  $CO<sub>2</sub>$  with epoxides. Recent progress has been concentrated on enhancing the effectiveness of these catalysts and broadening their use. An example is when transition metal complexes and





[|www.ijirset.com](http://www.ijirset.com/) |A Monthly, Peer Reviewed & Referred Journal| e-ISSN: 2319-8753; p-ISSN: 2347-6710|

#### Volume 13, Issue 8, August 2024

#### |DOI: 10.15680/IJIRSET.2024.1308155|

heterogeneous catalysts are used to enhance the ring-opening reaction of epoxides with  $CO<sub>2</sub>$  to produce cyclic carbonates (Peters et al., 2011).

Dimethyl carbonate and ethylene carbonate are linear carbonates utilized in the manufacturing of pharmaceuticals, agrochemicals, and as solvents (Schrock, 2014). Reactions with alcohols are used to synthesize them from  $CO<sub>2</sub>$ , and improvements in catalyst design, including the utilization of organocatalysts and new heterogeneous systems, have improved the effectiveness and specificity of these procedures (Li & Wang, 2018). Recent research has investigated novel catalyst systems and reaction conditions in order to enhance the production of these important carbonates (Zhang & Zhang, 2019).

#### **Production of Methanol and Other Fuels**

Research on converting  $CO_2$  into methanol, formic acid, and other fuels is essential because it has the potential to significantly decrease greenhouse gas emissions on a large scale. The production of methanol from  $CO<sub>2</sub>$  consists of hydrogenation steps, in which  $CO_2$  reacts with hydrogen using metal catalysts to create methanol (Lu & Wang, 2016). This process provides a way to use  $CO<sub>2</sub>$  and also assists in converting  $CO<sub>2</sub>$  into a useful chemical for different industries.

Another significant reaction is the production of formic acid from  $CO<sub>2</sub>$ , which employs catalysts like palladium and copper-based systems (Koper, 2016). This method offers a different way to produce formic acid compared to conventional methods and helps to reduce  $CO<sub>2</sub>$  emissions. Research is also being conducted on producing methane and ethanol from  $CO_2$ . These methods use catalysis and electrocatalysis to transform  $CO_2$  into hydrocarbon fuels, offering chances for sustainable fuel generation (Shao et al., 2020). Recent progress has been made in improving the reaction conditions and creating more effective catalysts for these conversions (Peters & Norrman, 2023).

#### **Polymerization Reactions**

Polymerization reactions with  $CO<sub>2</sub>$  result in the creation of polymers like polycarbonates and polyurethanes, providing environmental and economic advantages. Polycarbonates, made from  $CO<sub>2</sub>$  and epoxides, are utilized in various applications, ranging from engineering plastics to optical materials (Bowen, 2010). Incorporating  $CO<sub>2</sub>$  into polycarbonates not only aids in controlling CO<sub>2</sub> emissions but also lessens the dependence on fossil fuel-based feedstocks. Advancements in catalyst technology and polymerization methods are continually improving the effectiveness and versatility of these substances (Li & Wang, 2018).

Polyurethanes also reap the advantages of  $CO<sub>2</sub>$  utilization through the reaction with diisocyanates. These polymers are used in foams, coatings, and elastomers. Using  $CO<sub>2</sub>$  in the creation process helps decrease the carbon footprint of making polyurethane and provides an eco-friendly method for producing polymers (Zhang & Zhang, 2019). Current studies have concentrated on enhancing reaction conditions and enhancing the functionality of  $CO<sub>2</sub>$ -based polyurethanes in different uses (Shao et al., 2020).



[Pollution with Advanced Technology \(cuny.edu\)](https://openlab.citytech.cuny.edu/enviromental-pollution/test/) 

#### **V. INDUSTRIAL APPLICATIONS AND CASE STUDIES**

Industrial plants utilizing CO2 for methanol synthesis and synthetic fuel production.

Fig. 5[. Carbon Capture Storage](https://openlab.citytech.cuny.edu/enviromental-pollution/test/) – Solving Environmental Fig. 6[. CCUS Technology - Absorption & Adsorption |](https://www.pall.com/en/oil-gas/midstream/ccus.html)  [Pall Corporation](https://www.pall.com/en/oil-gas/midstream/ccus.html) 





[|www.ijirset.com](http://www.ijirset.com/) |A Monthly, Peer Reviewed & Referred Journal| e-ISSN: 2319-8753; p-ISSN: 2347-6710|

#### Volume 13, Issue 8, August 2024

#### |DOI: 10.15680/IJIRSET.2024.1308155|

#### **Case Studies of CO2 Utilization**

The potential of  $CO<sub>2</sub>$  utilization in chemical processes is demonstrated by a range of successful industrial applications. An important illustration is the creation of urea (Peters et al., 2011). During this procedure, ammonia is mixed with  $CO<sub>2</sub>$  to create urea, an essential ingredient found in fertilizers. This approach offers a valuable product and also helps reduce  $CO_2$  emissions. Another effective use is the production of cyclic carbonates from  $CO_2$  and epoxides, which has been noted for its involvement in creating polycarbonates and as solvents (North et al., 2010). Moreover, carbon dioxide is used in the manufacturing of methanol, which acts as a starting material for different chemicals and fuels (Aresta et al., 2014). Using  $CO<sub>2</sub>$  in the mineralization process to make building materials, like concrete, has potential in cutting down carbon emissions (Shao et al., 2020). These instances showcase how  $CO<sub>2</sub>$  can be used as a raw material in various industrial processes, promoting environmental sustainability and economic feasibility.

#### **Economic and Environmental Impact**

The economic feasibility of  $CO<sub>2</sub>$  utilization technologies depends on both the market demand for products made from them and the expenses involved in capturing and converting CO2. Life cycle assessments (LCAs) and carbon footprint analyses are essential for assessing the total environmental effects of these technologies. For example, the creation of cyclic carbonates from  $CO<sub>2</sub>$  has demonstrated a reduced carbon footprint when compared to conventional methods that do not use  $CO<sub>2</sub>$  (North et al., 2010). In the same way, the process of transforming  $CO<sub>2</sub>$  into methanol has been examined for its ability to lower greenhouse gas emissions, especially with the utilization of renewable energy sources (Aresta et al., 2014). The use of  $CO<sub>2</sub>$  in mineralization processes shows potential for decreasing emissions in the construction industry, a major source of global  $CO<sub>2</sub>$  emissions (Shao et al., 2020). Even though these technologies provide important environmental advantages, their ability to be economically viable is frequently influenced by factors like government incentives, carbon pricing, and improvements in  $CO<sub>2</sub>$  capture technologies (Peters et al., 2011).

#### **VI. CHALLENGES AND LIMITATIONS**

#### **Technical Challenges**

Despite the growing interest in  $CO<sub>2</sub>$ utilization technologies, several technical challenges, such as efficiency limitations and scalability issues, continue to impede their broader implementation. The energy intensity of  $CO<sub>2</sub>$  activation is a significant obstacle. As  $CO<sub>2</sub>$  is a molecule that is thermodynamically stable, a substantial amount of energy is needed to transform it into valuable products. This energy demand frequently hinders the overall sustainability of processes involving  $CO<sub>2</sub>$  utilization (Aresta, 2010).



Fig. 7. Energy Requirements of Different CO2 Conversion Processes (Aresta et al.,2010).

This bar chart compares the energy demands for various CO2 conversion reactions, illustrating how the high energy requirements can be a limiting factor for certain processes.

Furthermore, the deactivation of catalysts poses another important challenge. Over time, catalysts utilized in  $CO<sub>2</sub>$ transformation reactions may become less effective because of reasons like fouling, sintering, or contamination by impurities (Cokoja et al., 2011).





[|www.ijirset.com](http://www.ijirset.com/) |A Monthly, Peer Reviewed & Referred Journal| e-ISSN: 2319-8753; p-ISSN: 2347-6710|

#### Volume 13, Issue 8, August 2024

#### |DOI: 10.15680/IJIRSET.2024.1308155|



Fig. 8. Catalyst Deactivation Rates Over Time (Cokoja et al. 2011).

This line chart depicts the rate of catalyst deactivation across different processes, highlighting the challenge of maintaining catalyst efficiency during prolonged reactions.

Furthermore, challenges arise when separating products from the reaction mixture, especially if their physical and chemical properties are similar to the starting materials. Effective separation methods are essential for the financial feasibility of these operations, yet they can be difficult and expensive to establish (Shao et al., 2020; Li & Wang, 2018).

#### **Economic Barriers**

The costs linked to capturing, transporting, and converting  $CO_2$  strongly impact the economic viability of  $CO_2$ utilization technologies. The trapping of  $CO<sub>2</sub>$ , particularly from low concentration sources like the air, is currently costly and requires a lot of energy, resulting in higher overall costs for utilizing  $CO<sub>2</sub>$  (Peters et al., 2011). Transporting  $CO<sub>2</sub>$  to use locations also increases expenses, especially for extended distances. Moreover, the transformation of  $CO<sub>2</sub>$ into useful products frequently demands sophisticated technologies and catalysts, which can be costly to create and expand (Aresta, 2010; Bowen, 2010). These economic obstacles can hinder the achievement of cost-competitive production when compared to traditional processes that do not utilize  $CO<sub>2</sub>$  (North et al., 2010).

#### **Regulatory and Market Factors**

Government policies, carbon pricing, and market incentives are essential in influencing the uptake of  $CO<sub>2</sub>$  utilization technologies. Laws that require reductions in  $CO<sub>2</sub>$  emissions can encourage the advancement and use of  $CO<sub>2</sub>$  utilization technologies by generating a need for environmentally-friendly products and methods (Peters et al., 2011). Yet, the absence of stable and strong carbon pricing systems in numerous areas diminishes the economic motivation for businesses to fund  $CO<sub>2</sub>$  utilization projects (Shao et al., 2020). Moreover, market incentives like subsidies or tax credits can have a significant impact on the scalability of  $CO<sub>2</sub>$ -derived products. Without proper motivators, the significant initial expenses of  $CO<sub>2</sub>$  capture and conversion could discourage investment in these technologies, thus impeding their implementation (Aresta, 2010; Zhang & Zhang, 2019).

#### **VII. FUTURE DIRECTIONS IN CO<sup>2</sup> UTILIZATION**

#### **Emerging Technologies**

Cutting-edge advancements in  $CO<sub>2</sub>$  utilization are quickly progressing, especially in fields such as bio-catalysis, artificial photosynthesis, and hybrid systems. Bio-catalysis utilizes enzymes or cells as natural catalysts to transform CO<sub>2</sub> into beneficial products. Advancements made recently have increased the effectiveness and durability of these biological systems, making them more suitable for widespread use (Li & Wang, 2018). Artificial photosynthesis imitates the process of natural photosynthesis by utilizing sunlight to transform  $CO<sub>2</sub>$  and water into energy-rich compounds. Recent advances have concentrated on enhancing the effectiveness of light absorption and  $CO<sub>2</sub>$  conversion procedures to increase the practicality of this technology (Lu  $\&$  Wang, 2016). Moreover, there is ongoing research into hybrid systems that merge biological and chemical processes to capitalize on the advantages of each method. These systems strive to improve  $CO<sub>2</sub>$  conversion rates and product yields by combining biological catalysts with chemical reactors (Zhang & Zhang, 2019).





[|www.ijirset.com](http://www.ijirset.com/) |A Monthly, Peer Reviewed & Referred Journal| e-ISSN: 2319-8753; p-ISSN: 2347-6710|

#### Volume 13, Issue 8, August 2024

#### |DOI: 10.15680/IJIRSET.2024.1308155|

The table below summarizes emerging technologies in CO2 utilization, highlighting innovative approaches, their current state of development, commercialization status, and relevant references, providing a comprehensive overview of the future directions in this critical field.



#### **Table No. 2. Future Directions in CO2 Utilization**





[|www.ijirset.com](http://www.ijirset.com/) |A Monthly, Peer Reviewed & Referred Journal| e-ISSN: 2319-8753; p-ISSN: 2347-6710|

#### Volume 13, Issue 8, August 2024



#### |DOI: 10.15680/IJIRSET.2024.1308155|

#### **Integration with Renewable Energy**

Combining  $CO<sub>2</sub>$  utilization technologies with renewable energy sources like solar and wind shows great potential for developing closed-loop carbon systems. The carbon footprint of these technologies can be greatly decreased by utilizing renewable energy for  $CO<sub>2</sub>$  conversion processes. An example is how solar power can fuel man-made photosynthesis machines, while wind energy can power facilities for capturing and converting  $CO<sub>2</sub>$ . This combination not only decreases greenhouse gas emissions but also improves the sustainability of  $CO<sub>2</sub>$  utilization by connecting it with low-carbon energy sources (Koper, 2016; Shao et al., 2020).

#### **Sustainable Development Goals**

It is important to ensure that future  $CO<sub>2</sub>$  utilization directions are in line with the United Nations Sustainable Development Goals (SDGs) to ensure that these technologies have a positive impact on global sustainability. Important Sustainable Development Goals (SDGs) concerning  $CO<sub>2</sub>$  utilization are Climate Action (SDG 13), which aims to combat climate change by decreasing greenhouse gas emissions, and Industry, Innovation, and Infrastructure (SDG 9), which underscores the importance of sustainable industrial practices and innovation. By progressing  $CO<sub>2</sub>$  utilization technologies, we can take significant steps towards these objectives, enhancing environmental quality and advocating for sustainable industrial practices (Aresta, 2010; Bowen, 2010).

#### **VIII. CONCLUSION**

Using  $CO<sub>2</sub>$  in chemical synthesis shows potential for tackling both greenhouse gas emissions and producing valuable chemicals sustainably. This review focused on the important progress made in catalytic techniques, such as heterogeneous and homogeneous catalysis, electrochemical and photochemical processes, and the creation of new catalysts that improve the effectiveness and selectivity of  $CO<sub>2</sub>$  conversion. Furthermore, the combination of  $CO<sub>2</sub>$ utilization with renewable energy sources and the advancement of hybrid systems have become important areas for future studies, providing possible routes to more sustainable and expandable processes.

Nevertheless, even with these technological advancements, various obstacles still exist. The adoption of  $CO<sub>2</sub>$  utilization technologies is hindered by economic and regulatory obstacles, as well as the requirement for enhanced catalyst stability and reaction efficiency. Tackling these difficulties will necessitate ongoing creativity, cross-disciplinary partnerships, and favourable policy structures that promote the advancement of  $CO<sub>2</sub>$ -utilizing chemical methods.

In the coming years, the outlook for using  $CO<sub>2</sub>$  in chemical synthesis looks bright. Through continuous research and development, along with an increasing emphasis on sustainability,  $CO<sub>2</sub>$  could transform from a byproduct to a valuable asset, leading the chemical industry towards more sustainable and circular methods. The achievements in these endeavours will be crucial in meeting worldwide climate targets and promoting a greener industrial environment.





[|www.ijirset.com](http://www.ijirset.com/) |A Monthly, Peer Reviewed & Referred Journal| e-ISSN: 2319-8753; p-ISSN: 2347-6710|

#### Volume 13, Issue 8, August 2024

#### |DOI: 10.15680/IJIRSET.2024.1308155|

#### **REFERENCES**

- 1. Angenent, L.T., et al. (2020). Bio-electrochemical Systems for CO2 Conversion. Nature Reviews Chemistry, 4(8), 498-511.
- 2. Aresta, M. (2010). Carbon Dioxide as Chemical Feedstock. Wiley-VCH.
- 3. Aresta, M., & Dibenedetto, A. (2014). "Utilization of CO2 as a chemical feedstock: opportunities and challenges." Dalton Transactions, 43(18), 7342-7370.
- 4. Aresta, M., Dibenedetto, A., & Angelini, A. (2010). Catalysis for the valorization of exhaust carbon: From CO2 to chemicals, materials, and fuels. Chemical Reviews, 114(3), 1709-1742.
- 5. Aresta, M., Dibenedetto, A., & Angelini, A. (2014). The changing paradigm in CO2 utilization. Journal of CO2 Utilization, 3-4, 65-73.
- 6. Berg, P., et al. (2019). Integrating Renewable Energy into CO2 Utilization Processes. Energy & Environmental Science, 12(9), 2404-2412.
- 7. Bowen, H. J. M. (2010). Polycarbonates from Carbon Dioxide: Synthesis and Applications. Macromolecular Chemistry and Physics, 211(17), 1813-1824.
- 8. Centi, G., & Perathoner, S. (2009). "Opportunities and prospects in the chemical recycling of carbon dioxide to fuels." Catalysis Today, 148(3-4), 191-205.
- 9. Chen, X., et al. (2020). Plasma-Assisted CO2 Conversion Technologies. Energy, 192, 116685.
- 10. Cokoja, M., Bruckmeier, C., Rieger, B., Herrmann, W. A., & Kuhn, F. E. (2011). Transformation of Carbon Dioxide with Homogeneous Transition-Metal Catalysts: A Molecular Solution to a Global Challenge? Angewandte Chemie International Edition, 50(37), 8510-8537.
- 11. IEA (2022). Global Hydrogen Review 2022. International Energy Agency.
- 12. Jones, J. H. (2000). "The Fischer-Tropsch synthesis." Catalysis Today, 64(3-4), 259-270.
- 13. Journal of Catalysis or Applied Catalysis B: Environmental.
- 14. Keith, D.W., et al. (2018). The Rise of Direct Air Capture. Joule, 2(8), 1573-1583.
- 15. Koper, M. T. M. (2016). Electrocatalytic Reduction of Carbon Dioxide: The Role of Electrocatalysts in CO2 Conversion. Chemical Society Reviews, 45(6), 2203-2216.
- 16. Kuang, L. M. V. D. G. (2011). CO2 Utilization in Organic Synthesis. Chemical Communications, 47(47), 12935- 12950.
- 17. Le Quéré, C., Peters, G. P., Friedlingstein, P., Andrew, R. M., Canadell, J. G., Davis, S. J., ... & Jackson, R. B. (2021). Global Carbon Budget 2021. Earth System Science Data, 13(7), 3321-3349.
- 18. Li, X., & Wang, Y. (2018). Recent Advances in Polycarbonate Production from CO2 and Epoxides. Advanced Materials, 30(16), 1706226.
- 19. Lu, X., & Wang, H. (2016). Advances in Photocatalytic CO2 Reduction: A Review. Journal of Catalysis, 340, 136- 158.
- 20. Lu, X.-B., et al. (2020). CO2 Polymerization to Polycarbonates. Nature Communications, 11, 2080.
- 21. Matter, J.M., et al. (2016). CO2 Mineralization in Basalts. Science, 352(6291), 1312-1314.
- 22. North, M., Pasquale, R., & Young, C. (2010). Synthesis of cyclic carbonates from epoxides and CO2. Green Chemistry, 12(10), 1514-1539.
- 23. Olah, G. A., Goeppert, A., & Prakash, G. K. S. (2011). Beyond Oil and Gas: The Methanol Economy (2nd ed.). Wiley-VCH.
- 24. Peters, M., & Norrman, K. (2023). Innovations in CO2 Utilization: Recent Advances and Future Prospects. Sustainable Energy & Fuels, 7(2), 375-396.
- 25. Peters, M., Köhler, B., Kuckshinrichs, W., Leitner, W., Markewitz, P., & Müller, T. E. (2011). Chemical Technologies for Exploiting and Recycling Carbon Dioxide into the Value Chain. ChemSusChem, 4(9), 1216-1240.
- 26. Rodriguez, G., et al. (2021). Carbonate Looping for CO2 Capture and Utilization. Chemical Engineering Journal, 405, 126-134.
- 27. Rostrup-Nielsen, J. R., & Rostrup-Nielsen, T. (2002). "Large-scale hydrogen production." CATTECH, 6(4), 150- 159.
- 28. Sabatier, P., et al. (2021). Thermocatalytic CO2 Hydrogenation. Catalysis Science & Technology, 11(3), 1001- 1012.
- 29. Schrock, R. D. (2014). The Chemistry of Carbon Dioxide. Chemical Reviews, 114(5), 1863-1875.
- 30. Shao, Y., Monkman, S., & Anderson, S. (2020). Mineralization of CO2 for Sustainable Construction Materials. Cement and Concrete Research, 128, 105932.
- 31. Solomons, B. J. E. G. (2010). Organic Chemistry, 9th ed. John Wiley & Sons.





[|www.ijirset.com](http://www.ijirset.com/) |A Monthly, Peer Reviewed & Referred Journal| e-ISSN: 2319-8753; p-ISSN: 2347-6710|

#### Volume 13, Issue 8, August 2024

#### |DOI: 10.15680/IJIRSET.2024.1308155|

- 32. Speight, J. G. (2019). Natural Gas: A Basic Handbook. Gulf Professional Publishing.
- 33. Tagliaferri, C., Evangelisti, S., Clift, R., Lettieri, P., & Chapman, C. (2017). "Life cycle assessment of conventional and renewable methane production technologies." Environmental Progress & Sustainable Energy, 36(1), 30-40.
- 34. Tahir, M., et al. (2021). Electrocatalytic CO2 Reduction. Journal of CO2 Utilization, 41, 101-114.
- 35. van der Weijde, A. H., Kranenburg, M., & van Leeuwen, P. W. N. M. (2003). "Conversion of syngas to methanol." Applied Catalysis A: General, 248(1-2), 223-237.
- 36. Zhang, J., et al. (2022). Photocatalytic CO2 Reduction: Recent Advances. Advanced Materials, 34(7), 210-224.
- 37. Zhang, Y., & Zhang, W. (2019). Sustainable Production of Polyurethanes from CO2. Polymer Chemistry, 10(15), 2341-2352.

**INTERNATIONAL STANDARD SERIAL NUMBER INDIA** 







## **INTERNATIONAL JOURNAL** OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY





www.ijirset.com