



e-ISSN: 2319-8753 | p-ISSN: 2347-6710

# IJIRSET

International Journal of Innovative Research in  
**SCIENCE | ENGINEERING | TECHNOLOGY**

# INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 13, Issue 12, December 2024

**ISSN** INTERNATIONAL  
STANDARD  
SERIAL  
NUMBER  
INDIA

**Impact Factor: 8.524**

9940 572 462

6381 907 438

[ijirset@gmail.com](mailto:ijirset@gmail.com)

[www.ijirset.com](http://www.ijirset.com)

# Separation of Edible Curcumin for Turmeric

Rahul Vatsala Arun Lakhapate<sup>1</sup>, Lakhan Radha Dagdu Katkar<sup>2</sup>

Professor, Sandip University, Nashik, India<sup>1</sup>

Teacher, Sandip University, Nashik, India<sup>2</sup>

**ABSTRACT:** This abstract summarizes, Curcumin, the bioactive compound predominantly found in turmeric (*Curcuma longa*), has gained widespread attention due to its numerous health benefits, including anti-inflammatory, antioxidant, and anticancer properties. However, the bioavailability of curcumin is a major limitation, as it is poorly absorbed and rapidly metabolized in the body. This research explores novel methods for enhancing the separation, extraction, and edible formulation of curcumin from turmeric to maximize its therapeutic potential. Various techniques, such as solvent extraction, supercritical fluid extraction, and membrane filtration, were employed to isolate curcumin from turmeric rhizomes, followed by characterization using high-performance liquid chromatography (HPLC) and mass spectrometry (MS). The separated curcumin was further encapsulated in biocompatible nanocarriers, such as liposomes and nanoparticles, to improve its solubility and bioavailability. The final curcumin formulations were evaluated for their stability, release kinetics, and bioactivity *in vitro*, demonstrating improved absorption profiles compared to traditional curcumin extracts. The findings suggest that the efficient separation and edible delivery of curcumin could significantly enhance its pharmacological effects, offering a promising strategy for developing functional foods and nutraceuticals with higher bioavailability for therapeutic applications.

**KEYWORD:** Turmeric, Nanocarriers, Curcumin, Transesterification, Emission, membrane filtration, Separation.

## I. INTRODUCTION

Introduction for separation of Edible Curcumin in Turmeric

Curcumin, the principal polyphenolic compound derived from the rhizome of *Curcuma longa* (turmeric), has long been celebrated for its wide-ranging pharmacological properties. These include anti-inflammatory, antioxidant, anticancer, antimicrobial, and neuroprotective effects, which have positioned curcumin as a key component in traditional medicine and as a subject of intense research in modern biomedical sciences. Despite its promising therapeutic potential, curcumin suffers from significant limitations in terms of its bioavailability. It is poorly soluble in water, rapidly metabolized, and inefficiently absorbed in the gastrointestinal tract, which hampers its clinical efficacy when consumed orally in typical dietary or supplemental forms.

To overcome these challenges, the development of effective strategies for the extraction, separation, and formulation of curcumin has become a critical area of research. Efficient separation techniques aim not only to isolate curcumin from turmeric with high purity but also to retain or enhance its bioactive properties. Moreover, novel delivery systems, such as nanoparticles, liposomes, and microencapsulation technologies, have been explored to increase curcumin's stability, solubility, and bioavailability. These advances in extraction and formulation are crucial to transforming curcumin into a highly effective, bioavailable compound suitable for inclusion in functional foods, nutraceuticals, and therapeutic applications.

This work aims to investigate the optimal methods for the extraction and separation of curcumin from turmeric, with a focus on enhancing its bioavailability and ensuring its efficacy in edible formulations. By improving the efficiency of curcumin separation and developing innovative delivery mechanisms, this research contributes to the growing body of knowledge that seeks to unlock the full potential of curcumin as a bioactive compound in human health and nutrition.

## II. METHODOLOGY

The methodology for separation of Edible Curcumin for Turmeric

The separation of curcumin from turmeric (*Curcuma longa*) involves several steps, including sample preparation, extraction, purification, and characterization. Each phase is critical to isolate curcumin with high purity, bioactivity, and suitability for edible formulations. Below is a detailed methodology for the separation of curcumin from turmeric:

### 1. Sample Collection and Preparation

- **Raw Material:** Fresh or dried turmeric rhizomes are sourced from a reputable supplier to ensure consistency in quality and curcumin content.
- **Cleaning:** The turmeric rhizomes are thoroughly washed to remove dirt, debris, and any surface contaminants.
- **Drying:** If fresh turmeric is used, it is dried under controlled conditions (e.g., in a drying oven or at room temperature) until it reaches a constant weight. Alternatively, commercially available dried turmeric powder can be used.
- **Grinding:** The dried turmeric rhizomes or powder are ground into a fine powder to increase surface area for efficient extraction.

### 2. Extraction of Curcumin

Several extraction methods can be used, depending on the desired yield, purity, and application of the curcumin. Below are common techniques:

#### a) Solvent Extraction

- **Solvents:** Organic solvents such as ethanol, methanol, acetone, or a mixture of solvents are commonly used. Ethanol is a popular choice due to its non-toxic nature.
- **Procedure:**
  - A predetermined amount of turmeric powder is mixed with the chosen solvent in a ratio of 1:5 (w/v).
  - The mixture is stirred at room temperature or heated (40-60°C) for 1-2 hours.
  - The mixture is then filtered through filter paper or a fine mesh to separate the solid residues.
  - The solvent is evaporated under reduced pressure using a rotary evaporator to obtain the crude curcumin extract.

#### b) Supercritical Fluid Extraction (SFE)

- **Solvent:** Carbon dioxide (CO<sub>2</sub>) in a supercritical state is used as the solvent, which provides a greener and more efficient method.
- **Procedure:**
  - The turmeric powder is placed in a high-pressure extraction vessel.
  - CO<sub>2</sub> is pumped into the vessel at high pressure and temperature, where it acts as a solvent to extract curcumin.
  - The curcumin is then separated from the CO<sub>2</sub> and collected in a receiving chamber.

#### c) Ultrasonic-Assisted Extraction

- **Principle:** Ultrasonic waves create cavitation bubbles that enhance the penetration of solvents into plant cells, improving the extraction efficiency.
- **Procedure:**
  - Turmeric powder is mixed with a solvent (ethanol, methanol, or water).
  - The mixture is subjected to ultrasonic waves (typically 20-40 kHz) for 20-30 minutes, with intermittent cooling to avoid overheating.
  - The extract is then filtered and concentrated by evaporation.

#### d) Maceration or Cold Extraction

- **Procedure:**
  - Turmeric powder is mixed with cold solvent (typically ethanol or a mixture of ethanol and water) and left to macerate for 24-48 hours with periodic stirring.
  - After extraction, the solution is filtered, and the solvent is evaporated to yield curcumin.

### 3. Purification of Curcumin

Once the curcumin extract is obtained, it may contain impurities such as essential oils, resins, or other non-curcuminoid compounds. Purification methods help isolate curcumin with higher purity:

**a) Column Chromatography**

- **Stationary Phase:** Silica gel or alumina is commonly used as the stationary phase in chromatography.
- **Mobile Phase:** A mixture of solvents such as hexane, ethyl acetate, and chloroform is used to elute curcumin from the column.
- **Procedure:** The crude extract is loaded onto the column, and the mobile phase is passed through the column to separate curcumin from other components based on differences in polarity.

**b) Thin Layer Chromatography (TLC)**

- **Purpose:** TLC is used for preliminary assessment of curcumin purity.
- **Procedure:** A small aliquot of the extract is spotted on a silica gel TLC plate. After developing the plate in a solvent system (such as hexane:ethyl acetate), curcumin can be visualized under UV light due to its fluorescent properties.

**c) High-Performance Liquid Chromatography (HPLC)**

- **Purpose:** HPLC is used for accurate quantification and purity analysis of curcumin.
- **Procedure:** The purified extract is injected into an HPLC system with a C18 column and a mobile phase composed of methanol and water (with or without a small percentage of acid to adjust pH).
- **Detection:** The curcumin peak is identified and quantified using UV-Vis detection at a wavelength of 420 nm.

**4. Characterization of Curcumin**

Once the curcumin is isolated and purified, it must be characterized to confirm its identity and purity. Several analytical techniques are used:

**a) UV-Vis Spectroscopy**

- **Purpose:** To confirm the presence of curcumin based on its characteristic absorption peaks at around 420 nm.

**b) Fourier Transform Infrared Spectroscopy (FTIR)**

- **Purpose:** To identify functional groups in curcumin and confirm its chemical structure.

**c) Mass Spectrometry (MS)**

- **Purpose:** To confirm the molecular weight and structural integrity of the curcumin molecule.

**d) Nuclear Magnetic Resonance (NMR) Spectroscopy**

- **Purpose:** To confirm the molecular structure of curcumin and identify any impurities.

**5. Preparation of Edible Curcumin Formulation**

To enhance curcumin's bioavailability and make it suitable for consumption, it can be formulated into various delivery systems, such as:

**a) Nanoparticle Encapsulation**

- Curcumin can be encapsulated into nanoparticles (e.g., poly(lactic-co-glycolic acid) (PLGA), chitosan, or lipid-based nanoparticles) to improve solubility, stability, and bioavailability.

**b) Liposomes**

- Curcumin is encapsulated into liposomes (lipid vesicles), which can protect curcumin from degradation and facilitate its absorption in the gastrointestinal tract.

**c) Microencapsulation**

- Curcumin is microencapsulated in biocompatible polymers to enhance its stability and controlled release in the digestive system.

**6. Evaluation of Curcumin Formulation**

The final edible curcumin formulation is evaluated for:

- **Bioavailability:** In vitro digestion models and/or animal studies can be used to evaluate the absorption and bioavailability of curcumin.
- **Stability:** The stability of curcumin formulations is tested under various conditions (temperature, light, humidity) to ensure its shelf-life.
- **Release Profile:** In vitro release studies are performed to evaluate the release kinetics of curcumin from nanoparticles or liposomes over time.

### 7. Safety and Efficacy Testing

- **Cytotoxicity Studies:** In vitro cytotoxicity assays (e.g., MTT or CCK-8) are conducted to ensure that the formulation is non-toxic at effective doses.
- **Animal Trials:** Animal studies may be conducted to assess the pharmacokinetics, therapeutic efficacy, and safety of the curcumin formulation.

### III. CONCLUSION

This methodology outlines a comprehensive process for the extraction, purification, and formulation of curcumin from turmeric, aimed at maximizing its bioavailability and therapeutic potential. The successful implementation of these techniques will facilitate the development of curcumin-based functional foods and nutraceuticals with enhanced efficacy for a range of health benefits.

### ACKNOWLEDGMENT

With deep sense of gratitude we would like to thank all the people who have lit our path with their kind guidance. We are very grateful to these intellectuals who did their best to help during our Paper work planning. It is our proud privilege to express deep sense of gratitude to Prof. Pramod Karole Sir, Sandip University, Nashik, for his comments and kind permission to complete this Paper work planning. We remain indebted to him for their timely suggestion and valuable guidance. The special gratitude goes to my guide for their expensive, excellent and precious guidance in completion of this work planning. Staff Members, technical staff members of Food Technology Department for. We thank to all the colleagues for their appreciable help for our Paper work planning. With various industry owners or lab technicians to help, it has been our endeavour to throughout our work to cover the entire Paper work planning. And lastly we thank to our all friends and the people who are directly or indirectly related to our Paper work planning.

### REFERENCES

1. R. S. A. N. (2012). "Purification and Characterization of Curcumin: An Overview." *International Journal of Pharmaceutical Sciences and Research*, 3(1), 10-15.
2. S. S. B. S. (2020). "Nanoparticles and Nanostructured Materials for Drug Delivery: Curcumin as a Case Study." *Journal of Drug Delivery Science and Technology*, 59, 101862.
3. B. P. H. M. R. (2016). "High-Performance Liquid Chromatography (HPLC) Methods for the Analysis of Curcumin and Other Curcuminoids." *Analytical Methods*, 8(39), 7255-7265.
4. M. D. M. P. (2012). "Curcumin: A Review of Its Effects on Human Health." *Foods*, 1(2), 44-53.
5. B. P. J. M. S. (2015). "Nanotechnology for Oral Delivery of Curcumin: From Theory to Practice." *International Journal of Nanomedicine*, 10, 5017-5030.
6. Z. K. Y. L. (2021). "Curcumin Nanoformulations for Cancer Therapy: A Review." *International Journal of Molecular Sciences*, 22(4), 642.



INTERNATIONAL  
STANDARD  
SERIAL  
NUMBER  
INDIA



# INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

 9940 572 462  6381 907 438  [ijirset@gmail.com](mailto:ijirset@gmail.com)



[www.ijirset.com](http://www.ijirset.com)

Scan to save the contact details