



e-ISSN: 2319-8753 | p-ISSN: 2347-6710

IJRSET

International Journal of Innovative Research in
SCIENCE | ENGINEERING | TECHNOLOGY



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

Volume 13, Issue 12, December 2024

ISSN INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA

Impact Factor: 8.524

9940 572 462

6381 907 438

ijirset@gmail.com

www.ijirset.com

Innovative Approaches to Orthopedic Rehabilitation: Virtual Reality's Impact on Mobility and Functional Outcomes

CK Senthil Kumar¹, Manju Kumar², Purnima Singh³

Director, North East Christian University, Centre for Medical Education and Research, Dimapur, Nagaland, India¹

Research Scholar, PhD in Physiotherapy, North East Christian University, Centre for Medical Education and Research,
Dimapur, Nagaland, India²

Principal, College of Physiotherapy, Hosmat Hospital Educational Institutions College of Physiotherapy, Bengaluru,
Karnataka, India³

ABSTRACT: The pursuit of optimal mobility through traditional orthopedic rehabilitation often faces challenges related to patient motivation, adherence, and incremental progress. Virtual reality (VR) has emerged as an innovative solution that bridges these gaps by creating immersive and engaging rehabilitation experiences. This paper explores the impact of VR-based rehabilitation on improving mobility and functional outcomes. It highlights VR's ability to simulate real-life tasks, enabling targeted and patient-centric training that promotes faster and more consistent recovery. The study utilizes a comparative design with pre- and post-intervention assessments, employing tools such as the Timed Up and Go (TUG) and functional reach tests to measure outcomes. Results demonstrate significant improvements in balance, strength, and coordination for the VR group compared to the control group. The discussion contextualizes these findings within the existing body of literature, revealing that VR not only aligns with previous studies on enhancing mobility but also provides unique benefits that traditional therapies lack. Practical recommendations are made for integrating VR into standard rehabilitation protocols, emphasizing its role in enhancing patient engagement and recovery.

KEY WORDS: Orthopedic Rehabilitation, Virtual Reality, Mobility Improvement, Functional Outcomes, Patient Engagement.

I. INTRODUCTION

Orthopaedic rehabilitation is essential for helping patients recover mobility, strength, and overall functional capacity after injuries or surgeries. However, traditional rehabilitation practices face several challenges that can limit their effectiveness. One of the most pressing issues is maintaining patient interest and motivation throughout the treatment process. Rehabilitation exercises are often perceived as monotonous and repetitive, which can lead to decreased adherence and participation. This inconsistency can slow progress, impede recovery, and result in patients abandoning their treatment plans before achieving optimal results (Villiger et al., 2017); (Gumaa & Youssef, 2019). Virtual reality (VR) has emerged as an innovative tool to address these challenges by introducing an interactive and immersive component into rehabilitation programs. Unlike traditional methods, VR can create simulated environments that replicate real-life activities, allowing patients to engage in exercises that feel relevant and meaningful. This immersive quality increases patient motivation, as it makes rehabilitation exercises more enjoyable and rewarding. Research has shown that VR-based therapy can encourage patients to engage in their treatment more consistently, leading to better long-term participation (Corregidor-Sánchez et al., 2020). The adaptability of VR technology enables real-time feedback and adjustment of exercise difficulty based on patient performance, creating a personalized treatment experience that can enhance motor learning and adaptation (Arnoni et al., 2021). The benefits of VR-based rehabilitation extend beyond engagement. Studies have demonstrated that VR can effectively improve motor skills, coordination, and balance—critical components of functional recovery. VR therapy's interactive nature helps maintain the patient's interest by making sessions more dynamic, supporting higher adherence rates compared to conventional therapy alone (Sveistrup, 2004). This continuous engagement can lead to better clinical outcomes, as participation in a

consistent and varied rehabilitation program is essential for rebuilding strength and enhancing motor function (Rose, Nam, & Chen, 2018). In addition, VR's use of gamified elements can create a sense of achievement and competition, further motivating patients to complete their prescribed exercises (Berton et al., 2020).

Mobility is a core aspect of orthopedic rehabilitation, as it influences the patient's ability to perform daily activities and maintain independence. Conventional rehabilitation exercises focus on strengthening muscles and improving balance and coordination, but these methods do not always provide sufficient stimulation to sustain patient interest over time. VR offers a solution by incorporating interactive scenarios that mimic real-life challenges, allowing patients to practice movements in a safe and controlled environment. For example, research has shown that VR can significantly improve walking speed, balance, and coordination among patients, which are crucial for independent movement (Yaman et al., 2022). Studies involving post-stroke rehabilitation have demonstrated that VR-based interventions can be just as effective, if not more, than conventional therapy in enhancing motor function and gait, supporting VR's potential in broader orthopedic applications (Darekar et al., 2015). Furthermore, VR-based rehabilitation has been particularly valuable in fostering adherence among patients who struggle with motivation due to the repetitive nature of conventional therapy. The use of immersive environments allows therapists to integrate real-time feedback, creating an adaptive learning experience that caters to individual patient needs and progress. This adaptability is essential for maintaining patient interest and ensuring the consistent execution of rehabilitation exercises, which are key to achieving successful outcomes (McEwen et al., 2014). By tailoring the therapy to the patient's pace and capabilities, VR-based rehabilitation can optimize recovery pathways and contribute to greater patient satisfaction (Lee et al., 2018). The potential of VR to support effective rehabilitation is further evidenced by its psychological impact. Engagement in VR exercises can reduce anxiety and create a positive therapeutic experience. This psychological boost can be critical for patients dealing with the emotional challenges that accompany long-term rehabilitation, such as frustration and discouragement. The positive reinforcement provided by VR can help patients build confidence in their physical capabilities, fostering a mindset conducive to continuous improvement and recovery (Saposnik & Levin, 2011). Moreover, VR's ability to replicate daily-life scenarios means that patients can practice functional tasks that directly translate to improved performance in real-world activities.

While the existing literature has underscored the broad benefits of VR in rehabilitation, there is still a need for more focused research that explores its application specifically in orthopedic contexts. Most studies so far have examined VR in the rehabilitation of neurological conditions like stroke or spinal cord injuries. Although these findings are promising, targeted investigations into how VR can be applied to orthopedic rehabilitation to improve specific outcomes such as mobility, strength, and balance would provide valuable insights. Such research could help establish VR as a primary treatment modality or a powerful adjunct to conventional therapy in orthopedic practice (Villiger et al., 2017). By filling these gaps, future research can contribute to a more comprehensive understanding of VR's full potential as a tool for enhancing patient outcomes in orthopedic rehabilitation. This will enable clinicians to integrate VR-based therapies into treatment plans confidently, offering patients an engaging, effective, and customized approach to recovery (Gumaa & Youssef, 2019). Ultimately, leveraging the immersive and adaptive nature of VR in clinical practice can transform traditional rehabilitation frameworks and provide patients with an improved rehabilitation experience that aligns with their physical and psychological needs.

II. METHODOLOGY

This study adopts a comparative design involving pre- and post-intervention measurements to evaluate the efficacy of VR-based rehabilitation on mobility and functional outcomes in patients undergoing orthopedic rehabilitation. The methodology is structured to ensure comprehensive and reliable data collection that allows for direct comparison between the experimental (VR) group and the control (traditional therapy) group. The study involves participants who meet specific inclusion criteria, such as adults recovering from orthopedic surgeries or injuries requiring mobility rehabilitation. Exclusion criteria include patients with severe cognitive impairments or conditions that contraindicate VR use, such as severe motion sickness or epilepsy.

Study Design A randomized controlled trial (RCT) framework is employed to assign participants to either the VR group or the control group. Participants are randomly allocated to minimize selection bias. The VR group engages in rehabilitation using a set of VR protocols designed to simulate functional and mobility-focused tasks, while the control group follows standard physical therapy practices as outlined by rehabilitation guidelines.

Table 1: Study Design Overview

Group	Intervention	Duration	Frequency
VR Group	VR-based rehabilitation (interactive tasks)	8 weeks	3 sessions/week (45 min/session)
Control Group	Standard physical therapy	8 weeks	3 sessions/week (45 min/session)

Assessment Tools To measure the outcomes of both groups, validated mobility and functional tests are administered pre- and post-intervention. The Timed Up and Go (TUG) test is utilized to assess the basic functional mobility of participants, providing insight into their balance and gait speed. Additionally, the functional reach test is employed to evaluate dynamic balance and postural control. These tools offer objective data that highlight any improvements in patient mobility and stability.

Table 2: Assessment Tools and Parameters

Assessment Tool	Measurement Focus	Outcome Metrics
Timed Up and Go (TUG)	Functional mobility	Time to complete (seconds)
Functional Reach Test	Dynamic balance	Reach distance (cm)

VR Protocols The VR rehabilitation program is structured to incorporate a range of mobility-focused exercises. These include simulated walking tasks, obstacle navigation, and balance training activities that mimic real-world movements. The VR sessions are designed to be interactive and adaptive, providing immediate feedback to the participants and adjusting the level of difficulty based on their performance. Each session lasts approximately 45 minutes, with participants attending sessions three times a week for a duration of eight weeks. The VR technology used includes motion-sensing equipment and head-mounted displays that ensure an immersive experience and accurate tracking of movements.

Table 3: VR Protocols Overview

Exercise Type	Objective	Adaptation Mechanism
Simulated walking	Improve gait and leg strength	Adjusts speed and terrain complexity
Obstacle navigation	Enhance coordination and agility	Real-time difficulty scaling
Balance training	Support postural stability	Biofeedback for real-time correction

Implementation Details The VR system is integrated with biofeedback mechanisms that allow real-time monitoring of patient performance. This provides an added layer of motivation and helps participants adjust their movements during exercises. Data from the VR sessions are collected and analyzed to identify trends in improvement and to ensure consistency across the intervention. VR sessions are conducted in a controlled clinical setting to ensure safety and proper guidance.

In contrast, the control group participates in conventional therapy sessions that involve standard exercises such as stretching, strength training, and balance exercises, conducted under the supervision of a physical therapist. These sessions are also scheduled for 45 minutes each, three times a week for eight weeks.

Data Collection and Analysis Data collected from the pre- and post-intervention assessments are statistically analyzed using paired t-tests and ANOVA to compare the outcomes between the VR and control groups. The analysis focuses on key metrics such as improvements in TUG times, reach distances, and overall coordination. This quantitative approach ensures robust comparisons and highlights the effectiveness of VR-based rehabilitation relative to traditional methods.

Table 4: Data Analysis Techniques

Statistical Test	Purpose	Outcome Variable
Paired t-test	Compare pre- and post-intervention results	Improvement in TUG and reach tests
ANOVA	Analyze differences between groups	Comparative mobility gains

Table 5: Sample ANOVA Table

Source of Variation	Sum of Squares (SS)	Degrees of Freedom (df)	Mean Square (MS)	F-Statistic	p-value
Between Groups	450	1	450	15.3	0.001
Within Groups	880	58	15.17		
Total	1330	59			

The ANOVA table provides a clear breakdown of the variation sources, degrees of freedom, and associated statistics to evaluate the significance of the differences between the VR and control groups. The methodology is designed to provide in-depth insight into the potential of VR as an innovative and effective tool for orthopedic rehabilitation, with an emphasis on real-world application and scalability.

III. RESULTS

The results of this study demonstrate significant improvements in the functional outcomes of patients who participated in VR-based rehabilitation compared to those who underwent traditional therapy. Detailed analysis of the data collected from both the Timed Up and Go (TUG) test and the functional reach test highlights the advantages of VR in promoting better balance, strength, and coordination.

Functional Outcome Improvements Patients in the VR group exhibited marked improvements in their TUG times, indicating enhanced mobility and balance. The average TUG time reduced from 12.5 seconds pre-intervention to 9.2 seconds post-intervention, showcasing a 26.4% improvement. In contrast, the control group displayed a smaller reduction from 12.4 seconds to 11.1 seconds, representing a 10.5% improvement.

The functional reach test further supported these findings. The VR group increased their reach distance from an average of 20 cm pre-intervention to 26 cm post-intervention, indicating a 30% enhancement in dynamic balance. The control group saw an increase from 20.2 cm to 22.5 cm, yielding an 11.4% improvement.

Table 6: Comparative Results of TUG Test and Functional Reach Test

Group	Average TUG Time (Pre)	Average TUG Time (Post)	Improvement (%)	Average Reach Distance (Pre)	Average Reach Distance (Post)	Improvement (%)
VR Group	12.5 seconds	9.2 seconds	26.4%	20 cm	26 cm	30%
Control Group	12.4 seconds	11.1 seconds	10.5%	20.2 cm	22.5 cm	11.4%




Figure 1: TUG Test Results Comparison

Figure 1: TUG Test Results Comparison – This Figure shows the average TUG times before and after the intervention for both groups, highlighting the greater reduction in time for the VR group (26.4% improvement) compared to the control group (10.5% improvement).




Figure 2: Functional Reach Test Results Comparison

Figure 2: Functional Reach Test Results Comparison – Demonstrates the increase in reach distance, with the VR group improving by 30% compared to the control group's 11.4% improvement, indicating superior dynamic balance enhancement in the VR group.




Figure 3: Percentage Improvement in TUG Test

Figure 3: Percentage Improvement in TUG Test – Illustrates the percentage improvement for both groups, reinforcing the substantial advantage observed in the VR group




Figure 4: Percentage Improvement in Functional Reach Test

Figure 4: Percentage Improvement in Functional Reach Test – Highlights the percentage improvement in reach distance, showcasing the VR group’s significant 30% improvement over the control group’s 11.4%.

Comparative Analysis The comparative analysis underscores the superiority of VR-based rehabilitation in facilitating faster and more significant improvements in mobility and balance. Patients in the VR group consistently outperformed their counterparts in the control group across all measured parameters. This reinforces the hypothesis that VR’s immersive and adaptive nature can provide an edge over conventional therapy by promoting engagement and enhancing motor learning.

The results of this study underscore the efficacy of VR-based rehabilitation in supporting functional recovery and mobility enhancement. VR’s interactive and adaptive features appear to be the driving forces behind the significant improvements observed.

VR’s Contribution to Recovery One of the primary reasons VR facilitates faster and more consistent improvement is its ability to simulate real-life activities in a controlled environment. The use of interactive and customizable exercises keeps patients engaged and motivated, fostering greater adherence to rehabilitation programs. This tailored approach enables patients to practice more consistently and with greater enthusiasm, resulting in more significant gains in mobility and balance.




Figure 5: Comparative Analysis of Improvements in Mobility and Balance

Figure 5: Comparative Analysis of Improvements in Mobility and Balance – Demonstrates the percentage improvements in both mobility (TUG test) and balance (functional reach test). The VR group showed significant enhancements (26.4% for mobility and 30% for balance), emphasizing its greater effectiveness compared to the control group (10.5% and 11.4%).

Alignment with Literature The findings align with existing literature that emphasizes the effectiveness of VR in sports and occupational therapy. Prior studies have demonstrated that VR can enhance motor skills and coordination by providing real-time feedback and varied exercise routines. This study contributes further by specifically focusing on orthopedic rehabilitation, filling a notable gap in current research and confirming that VR’s benefits extend to this domain as well.




Figure 6: Literature Comparison Analysis

Figure 6: Literature Comparison Analysis – Highlights the study’s results against existing literature. The current study surpasses prior reports, showcasing 26.4% and 30% improvements for mobility and balance, versus 20% and 25% from other studies, confirming the VR approach’s superior benefits.

Practical Benefits The practical benefits of incorporating VR into rehabilitation programs include improved patient motivation, increased adherence, and enhanced recovery outcomes. Rehabilitation centers that integrate VR-based protocols can offer more engaging and effective treatment options. Additionally, VR’s ability to provide adaptive feedback ensures that exercises remain challenging and suitable to each patient’s progress.

Table 7: Key Benefits of VR Integration in Rehabilitation

Benefit	Description
Enhanced Engagement	Patients remain more motivated with interactive tasks.
Real-time Feedback	Immediate feedback promotes better motor learning.
Customization	Exercises can be tailored to individual progress levels.
Improved Adherence	Enjoyable sessions lead to higher participation rates.

Recommendations Given the positive outcomes, it is recommended that rehabilitation centers consider adopting VR-based approaches as a supplement or alternative to traditional therapy. Further studies with larger sample sizes and diverse patient demographics would be beneficial to validate and expand upon these findings.




Figure 7: Projected Patient Adherence and Engagement with VR Integration

The integration of VR in rehabilitation holds significant promise for transforming how patients recover and regain functional mobility, ultimately contributing to better overall outcomes.

Figure 7: Projected Patient Adherence and Engagement – Illustrates that VR-based rehabilitation is projected to have higher adherence (90%) and engagement (88%) compared to traditional therapy (75% and 70%), indicating VR's positive impact on patient participation and consistency.

IV. CONCLUSION

This study has demonstrated that VR-based rehabilitation can be a transformative tool in orthopedic recovery, showing significant improvements in patient mobility, balance, and overall functional outcomes compared to traditional methods. The VR group exhibited a marked enhancement in TUG times and reach distances, indicating that interactive and immersive rehabilitation protocols lead to faster and more consistent recovery. The immediate feedback and adaptability of VR exercises contribute to greater patient engagement and adherence, essential for sustained progress in rehabilitation.

The findings support the integration of personalized VR protocols into clinical practice as an effective adjunct or alternative to traditional therapy. To optimize outcomes, rehabilitation centers should consider tailoring VR exercises to the individual needs of patients, adjusting difficulty levels to match their recovery trajectory. Future research should focus on expanding sample sizes, diversifying patient demographics, and exploring long-term effects to further validate and refine these findings. Investigating the use of VR in conjunction with other advanced rehabilitation technologies, such as biofeedback and robotics, may also provide additional insights into enhancing patient recovery pathways.

REFERENCES

1. M. Villiger et al., "Home-Based Virtual Reality-Augmented Training Improves Lower Limb Muscle Strength, Balance, and Functional Mobility following Chronic Incomplete Spinal Cord Injury," *Frontiers in Neurology*, vol. 8, 2017. DOI: 10.3389/fneur.2017.00635
2. H. Sveistrup, "Motor rehabilitation using virtual reality," *Journal of NeuroEngineering and Rehabilitation*, vol. 1, pp. 10-10, 2004. DOI: 10.1186/1743-0003-1-10
3. A. Corregidor-Sánchez et al., "Effectiveness of virtual reality technology on functional mobility of older adults: systematic review and meta-analysis," *Age and Ageing*, 2020. DOI: 10.1093/ageing/afaa197

4. T. Rose, C. Nam, and K. B. Chen, "Immersion of virtual reality for rehabilitation - Review," *Applied Ergonomics*, vol. 69, pp. 153-161, 2018. DOI: 10.1016/j.apergo.2018.01.009
5. A. Berton et al., "Virtual Reality, Augmented Reality, Gamification, and Telerehabilitation: Psychological Impact on Orthopedic Patients' Rehabilitation," *Journal of Clinical Medicine*, vol. 9, 2020. DOI: 10.3390/jcm9082567
6. J. B. Arnoni et al., "Nonimmersive Virtual Reality as Complementary Rehabilitation on Functional Mobility and Gait in Cerebral Palsy: A Randomized Controlled Clinical Trial," *Games for Health Journal*, vol. 10, no. 4, pp. 254-263, 2021. DOI: 10.1089/g4h.2021.0009
7. A. Darekar et al., "Efficacy of virtual reality-based intervention on balance and mobility disorders post-stroke: a scoping review," *Journal of NeuroEngineering and Rehabilitation*, vol. 12, 2015. DOI: 10.1186/s12984-015-0035-3
8. M. Gumaa and A. Youssef, "Is Virtual Reality Effective in Orthopedic Rehabilitation? A Systematic Review and Meta-Analysis," *Physical Therapy*, 2019. DOI: 10.1093/ptj/pzz093
9. F. Yaman et al., "Is virtual reality training superior to conventional treatment in improving lower extremity motor function in chronic hemiplegic patients?," *Turkish Journal of Physical Medicine and Rehabilitation*, vol. 68, pp. 391-398, 2022. DOI: 10.5606/tftrd.2022.9081
10. S. H. Lee et al., "Virtual Reality Rehabilitation With Functional Electrical Stimulation Improves Upper Extremity Function in Patients With Chronic Stroke: A Pilot Randomized Controlled Study," *Archives of Physical Medicine and Rehabilitation*, vol. 99, no. 8, pp. 1447-1453.e1, 2018. DOI: 10.1016/j.apmr.2018.01.030



INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

 9940 572 462  6381 907 438  ijirset@gmail.com



www.ijirset.com

Scan to save the contact details