



International Journal of Innovative Research in Science Engineering and Technology (IJIRSET)

(A Monthly, Peer Reviewed, Refereed, Scholarly Indexed, Open Access Journal)



Impact Factor: 8.699

Volume 14, Issue 4, April 2025

A Review on Evaluating the Nutritional Properties of Food

Hemalatha Sattu, P.Vivek Sagar, P.Sriharsha, P.Haripriya, K.Nikitha

Professor & HOD, Dept. of Pharmaceutical Chemistry, Sarojini Naidu Vanita Pharmacy Maha Vidyalaya, Secundrabad,
Telangana, India

Professor & HOD, Dept of Pharmaceutical Analysis, Sarojini Naidu Vanita Pharmacy Maha Vidyalaya, Secundrabad,
Telangana, India

M.Pharm, Sarojini Naidu Vanita Pharmacy Maha Vidyalaya, Secundrabad, Telangana, India

ABSTRACT: Many techniques or indicators are used to evaluate the nutritional content of the meal; However, a systematic assessment system is not available, and it is difficult to fully describe the overall nutritional value of the food with a single guide. Typical evaluation techniques and nutritional food content indicators are included in this method in this article. The goal of this assessment is to provide a process to assess the nutrients of meals and help scientists choose simpler and more effective evaluation techniques based on foods or relevant indicators. This process is to choose a three -angle evaluation technique that includes the entire scope of nutritional quality of a product that it is permanent to think reasonably in the fields of food, rural science and food

KEYWORDS: nutrition evaluation, nutritional value of food, evaluation techniques, indications for evaluation

I. INTRODUCTION

A food is a complex mixture of nutrients and other molecules interacting inside and between meals. Foods have a great impact on human metabolism, digestion and development. Food -related areas include nutrition, food science, agricultural science and many other areas. It is often realized that agriculture can help reduce malnutrition. A new agricultural model called nutrition agriculture promotes agricultural experts to focus more on the nutritional value of food. The expansion of specific treatment can be helped by nutrition sensitive agricultural initiatives. Find food with high nutritional value and use this information to determine their costs as a concept provided by agricultural economists. Food science researchers are mainly interested in treatment and quality technology. Nutrition content, qualitative quality, packaging, storage and storage. Consumers and experts have an unfavorable idea about meals processed due to their disagreements with food and health. Always ask questions if the food contribution of extremely - upfs) has a significant impact on the overall nutritional value of contemporary regimes. The main goal of science on nutrition is how nutrients affect human health. It broke up complicated nutritional needs into digestive proposals in the form of a food visa to help prevent disease. Many studies conducted after more than a century of development have revealed that food deficiency, such as incomplete consumption of nutrients, is the main cause of most chronic diseases. There are current problems . In the clinical context, the term "nutrition" is often used, the human body serves as an assessment object. Research on the evaluation of nutrients of most food tools, and different fields with different definitions and explanations of the same topic. For example, food scientists often use the type and number of nutrients in the product to determine its nutritional value. However, the genre and number of nutrients may not be enough to show the nutritional value of the food. In addition, nutrition experts and researchers emphasize how nutrients or the final food system affects human health. An evaluation technique with different methods should be used to evaluate the nutritional value

II. ASSESSMENT OF WHOLESOME ESTEEM

Creating a strong and deliberate system for assessing the dietary Choosing dependable files and criteria is the establishment of logical appraisal methods that are viable in evaluating the dietary substance of nourishment. Various assessment methods have been created and are habitually utilized. In arrange to compare the strategies and give modern

hypothetical direction or fortify the creation of unused strategies for deciding the wholesome esteem of nourishment, the most objective of this think about was to accumulate all of the strategies that have been utilized in this way distant, show them in a methodical way, and after that completely analyse them. As a result, we considered the contrasts within the assessment strategies, as well as their benefits and disadvantages. The evaluation strategies were categorized into the taking after bunches utilizing this assessment methodology.

Global Food Security Index, 2020
(Overall score 0–100, 100 = best environment)




A collection of files from 113 diverse countries make up the Worldwide Nourishment Security Record. It gages nourishment security within the larger part of the world's nations.[1] It was at first discharged in 2012, and The Economist's insights segment supervises and upgrades it each year. Assessment Strategy	Creator	Year	Definition	Computing Strategy	Components Assessment	quality list
AAS, or amino acid score	Bano et al.	1982	a rating that surveys how near an vital amino	AAS = (amino acid concentration in test protein (mg/g)) /	Amino acid substance in a test of protein Amino	Amino acid score

			corrosive within the protein beneath test is to its identical within the reference protein demonstrate.	(coordinating fundamental amino acid substance within the FAO or WHO scoring standard show (mg/g))	corrosive score		
PDCAAS stands for Protein-Digestibility-Corrected Amino Acid Score.	Eggum et al.	1991	A developing rodent was utilized as a show for the grown-up human, and the proportion of the amount of the first-limiting dietary basic amino corrosive within the protein source to the amino corrosive requests of a child matured 1-2 a long time was balanced for protein digestibility utilizing genuine fecal nitrogen digestibility.		Protein digestibility, a child's 1-2 year old's amino acid needs, and the quantity of first-limiting amino acids	PDCAAS score	
Score for Indispensable Digestible Amino Acids (DIAAS)	Wolfe et al.	2016	Based on the digestibility of essential amino acids (IAAs) and the amino acid demand pattern.	$\text{DIAAS (\%)} = 100 \times \frac{\text{(mg of digestible dietary IAA in 1 g of the dietary test protein)}}{\text{(mg of the same amino acid in 1 g of the reference protein)}}$	Histidine (His), isoleucine (Ile), leucine (Leu), valine (Val), lysine (Lys), threonine (Thr), phenylalanine (Phe), methionine (Met), and tryptophan (Trp) are among the IAAs' amounts and profiles.	DIAAS score	
Essential Amino Acid Index (EAAI).	Oser et al.	(1951)	A geometric cruel is utilized to compute an generally comparison of all fundamental amino acids within the protein beneath test to all essential amino acids within the reference protein demonstrate.		Each critical amino acid	EAAI	

The Index of Nutrition Quality (INQ)	Gholamalizadeh et al.; Sorenson et al.	(1976; 2021)	A technique that is particularly important for evaluating clinical nutritional concerns is the quantitative and qualitative analysis of individual foods, meals, and diets.	NQ is derived by dividing a nutrient's RDA in 1000 kcal by its amount in 1000 kcal of food or diet.	Illustrations incorporate thiamin, riboflavin, niacin, pantothenic, vitamin B6, folate, vitamin B12, copper, magnesium, zinc, calcium, selenium, vitamin A, vitamin C, vitamin E, and vitamin D.	INQ results
Naturally Rich in Nutrients (NNR)	Drewnowski et al.	2005	14 nutrients' crucial values (DVs) in 2000 kcal of nourishment.	NNR is rise to to $\sum \%DV_{2000 \text{ kcal}/14}$	Monounsaturated fat, potassium, zinc, calcium, press, protein, vitamins A and C, thiamine, riboflavin, vitamin B12, folate, vitamin D, and vitamin	NNR rating
Calories-for-Nutrients (CFN)	Drewnowski et al.	2005	The number of calories needed to get an extra 1% DV for many important nutrients.	$.ED / (\sum \%DV_{100 \text{ g}} / 13) = CFN$	Niacin, folic acid, magnesium, zinc, calcium, iron, vitamin A and C, thiamine, riboflavin, vitamin B6, vitamin B12, and protein	CFN rating
The Recommended to Restricted Food Score (RRR) ratio	Scheidt and associates	2004	Depending on the item name, the proportion of "great" to "terrible" supplements, and the item's vitality level	RRR is equal to $(\sum \%DV_{\text{recommended}} / 6) \text{ divided by } (\sum \%DV_{\text{restricted}} / 5)$.	Five supplements (vitality, immersed fat, cholesterol, sugar, and salt) had undesirable RRR evaluations, though six supplements (protein, calcium, press, vitamin A, vitamin C, and fiber) were respected positive a priori.	RRR Score
Index of Dietary Quality (DQI)	Patterson et al.	1994	An tool to quantify overall diet quality that shows a risk gradient for major food-related chronic illnesses based on US dietary guidelines from Diet and Health.	Cutoffs were created for index scoring, the Diet and Health recommendations were weighted, and scores were added together for all of the recommendations	salt, calcium, grains and legumes, fruits and vegetables, cholesterol, energy from fat, energy from saturated fat	DQI total score (0–16)

The updated Dietary Quality Index (DQI-R)	Haines et al.	1999	a 1994 overhaul to the DQI based on USDA information, which reflected the population's most later dietary proposals.	Ten focuses are included to the in general DQI-R score for each of the ten components	Dietary cholesterol (mg), calcium, RDA press every day, dietary assortment, dietary control, day by day suggested servings of natural products, vegetables, and grains, vitality from fat, vitality from soaked fat, and other variables	Add up to DQI-R score (0–100)
International Diet Quality Index (DQI-I)	Kim et al.	2003	A composite metric for food quality was developed to assess the nutritional value of a diet both inside a nation for monitoring reasons and internationally for comparison analysis.	Scores for each component are summarized in each of the four primary categories, and the scores for all four categories are summed	Adjust (macronutrient proportion, greasy corrosive proportion), differences (both generally and inside protein sources), ampleness (natural products, vegetables, grains, fiber, protein, press, calcium, and vitamin C), and control (add up to fat, immersed fat, cholesterol, salt, and empty-calorie suppers)	Total DQI-I score (0–100)
Glycemic Index (GI)	Jenkins et al.	1981	The glycemic effect of accessible carbs in diet relative to the effect of an equal amount of glucose.	GI is equal to $\frac{\text{Iaucfood/Iaucglucose}}{\text{Wt glucose/Wt food-available carbohydrate}} \times 100\%$	The impact of available carbohydrates on the GI score	GI score (less than 55, low GI; 55–69, medium GI; greater than 70, high GI) of human blood



A sensory lexicon is a list of words and definitions that describe the sensory qualities of a food or drink. It's a standardized vocabulary that's used to communicate with sensory panels, scientists, and other professionals in the food industry.

The benefits and drawbacks of various approaches

Types of Methods	Principal Methods	Advantages	Disadvantages	Examples
According to Food Nutrients	Comparison of Nutrients	Quick & easy	Unable to accurately depict the health benefits of eating Nutrient	comparison between food items and control samples or FCTs
	Chemical Scoring	Scientific and effective; further analysis of nutrient status	Individual scores can only reflect the nutritional characteristics of food from a certain angle; scores should be combined to reflect the nutritional properties of food	The basic amino acid list (EAAI), index of nutrition quality (INQ), naturally nutrient rich (NNR), digestible indispensable amino acid score (DIAAS), protein-digestibility-corrected amino acid score (PDCAAS), amino acid score (AAS), and calories-for-nutrients (CFN)
Using Perceptions from the Senses	An explanation of color, flavor, and scentintuitive	readily embraced by customers	It requires expert researchers to finish; Individual tastes differ greatly. Sensory assessment	Sensory assessment

S.NO	Functional Components	Components of Nutraceutical Foods:	Possible Advantages	Citation
1	whole grain, Garlic with Onion	FOS, Inulin	Obesity, diabetes, constipation, and chronic illness prevention and therapy.	(Megur and others, 2022).
2	Apple, tea, chocolate, and grapes	Catechins	Chronic illness prevention and therapy.	(Koushki and colleagues, 2018)
3	Citrus fruits, pumpkin, squash, and carrots	β -carotene	For healthy skin and mucous membranes, for a robust immune system, and for eye health.	(Carazo and others, 2021).
4	Red wine and grapes	resveratrol	Alzheimer's illness, diabetes, cancer, and antioxidants.	(Koushki and colleagues, 2018)
5	Rye, flax, onion, and garlic	Adenosine, lignan	Reduced risk of osteoporosis, breast cancer, menopausal symptoms, and heart disease	(Jang and others, 2022).
6	Wine in red	anthocyanates,	anti-inflammatory, anti-cancer, anti-microbial, anti-obesity, and anticipation of cardiovascular disorders	Ghoshal (2024)
7	The majority of plants	Cellulose	Blood sugar, cholesterol, and constipation	Haub and Lattmer (2010)
8	Psyllium husk	Psyllium	Reduce your cholesterol. Psyllium is used to treat hemorrhoids, irritable bowel syndrome, and other digestive issues. It can also help reduce diarrhea and constipation.	(Garg, 2021)
9	Dairy and Beef	The linoleic acid conjugate	antihypertensive, antidiabetic, anticarcinogenic, and anti-obesity qualities, as well as the ability to prevent metabolic syndromes and lifestyle disorders.	(Yanagita and Koba, 20)

TABLE 2 : EXAMPLES OF FUNCTIONAL FOOD COMPONENTS AND POTENTIAL BENEFITS

III. METHODS OF EVALUATION USING FOOD NUTRIENTS

1. The Nutrient Comparison Method

nourishments are differentiated with the values recorded within the food composition table (FCT). The food's introductory wholesome esteem may at that point be discovered utilizing the dietary reference intakes (DRIs) in agreement with the prescribed consumption. A clear strategy of controlling nourishment labeling, wellbeing claims, showcasing, and advancement is through nutrient profiling, which is the method of gathering or positioning things concurring to their dietary substance. Food composition tables and databases (FCT/FCDB) compile information on the vitality and dietary substance of foods in a certain country or range. Around 100 countries Since the 1980s, a number of FCDBs have developed, counting those for Europe, Oceania, North America, Asia, and Latin America. Both Eastern and Western Europe are included in Euro cuisine. The US Department of Agriculture's (USDA) nourishment composition data online is centralized at Food Data Central (FDC), which gives nourishment sustenance data for Americans. The Global Branded Foods Products Database (GBFPD) is one of five sorts of information given by the Food and Nutrient Database for Dietary Studies (FNDDS), standard reference legacies (SR Bequest), foundation foods (FF), and experimental foods (EF). The FDC offers reliable, web-based, reasonable, and effectively accessible data approximately the supplements and other fixings in dinners to fulfill the continually moving requests of different groups of onlookers. More than 600 items' food composition data (FCD), with up to 87 major components It has been a portion of the New Zealand Food Composition Database (NZFCDB) since 2010. To offer assistance customers get it the wholesome esteem of food and make educated eating choices, China's FCT gives common nutrition information on approximately 4710 crude plant and creature components. We too found that national tables of nourishment composition are seldom the same. It is challenging to compare dietary substance since numerous items have different portrayals or presentations.

2. Method of Chemical scoring

The primary component of living things, proteins are a type of organic macromolecules that make up cells' fundamental organic substance. Without protein, life would not exist. The building blocks of proteins are amino acids. As a result, protein quality has long been used to gauge a food's nutritional worth. Chemical scoring, which includes creating an suitable scientific show based on the wholesome substance, is generally utilized for protein appraisal. The degree to which an basic amino corrosive within the protein being tried is indistinguishable to its equivalent essential amino corrosive within the reference protein show is measured by the amino acid score (AAS). It may be a commonly utilized metric to survey the dietary substance of suppers tall in protein. The Food and Agriculture Organization of the United Nations (FAO) made the protein-digestibility-corrected amino acid score (PDCAAS) in 1990 as a way to evaluate the quality of proteins. A PDCAAS of 1.0 shows that all of the minimal needs for essential amino acid (IAA) intake would be fulfilled in the event that the sum of test protein devoured was identical to the estimated average requirement (EAR) for protein. The truncation of the PDCAAS is 1.0, indicating that it is unable to separate between the relative quality of diverse premium dietary proteins. As a result, Wolfe et al. and Mahtai et al. proposed that the digestible amino acid scores (DIAAS) are a stronger strategy than the PDCAAS for assessing the protein substance of meals. It is hypothetically conceivable to score all dietary proteins concurring to their quality since the DIAAS isn't abbreviated for a single-source protein just like the PDCAAS. A geometric cruel that compares all of the basic amino acids within the protein beneath test to all of the fundamental amino acids within the reference protein show is the essential amino acid index (EAAI), which is another sign. Protein does not whole up a food's nutritious qualities, indeed in case it may be a critical portion of it. Various markers of nutritional substance for entirety dinners have been made and put into utilize. Various measurements, counting nutrients-per-calorie and calorie-to-nutrient appraisals, have been made to assess the dietary thickness of nourishments. A instrument for both quantitative and subjective inquire about on certain foods, dinners, and diets is the food dietary quality list, often referred to as the file of nourishment quality, or INQ. It is especially accommodating for evaluating clinical dietary issues. The naturally nutrient-rich (NNR) score, which is decided by calculating the mean rate every day values (DV) for 14 supplements in 2000 kcal of nourishment, may be utilized to dole out supplement thickness appraisals for nourishments both inside and inside nourishment bunches. Nourishment instruction and dietary suggestions can be incredibly improved by the supplement thickness approach. The fetched in calories required to get an extra 1% DV for a assortment of important nutrients is known as the calories-for-nutrients (CFN) score, and it may be utilized to evaluate the coordinate relationship between dietary esteem and dietary vitality.

3. Evaluation Techniques

Based on Food Function. A developing sum of considers has appeared that a single or limited supplement assessment cannot accurately reflect a diet's wholesome composition. A dietary framework may have a diverse relationship with wellbeing markers than person components examined autonomously. Hence, from the viewpoint of adjust, differing qualities, ampleness, As dietary markers, control, count calories, and nourishment category quality have all been recommended. The study of disease transmission states that a food's dietary worth ought to take into consideration its positive impacts on the human body, such as the avoidance and treatment of diabetes, cardiovascular infection, and other constant ailments; its chemical components fair constitute its wholesome establishment. In 1994, Patterson et al. made a diet quality index (DQI) based on the Eat less and Wellbeing eating rules for Americans. Eight components made up the first DQI, and its values extended from (great count calories) to 16 (terrible slim down). Haines et al. overhauled the DQI in 1999 and produced the DQI-Revised (DQI-R), which consolidates pointers of dietary assortment and balance in conjunction with more exact nourishment parcel estimations. The DQI-International (DQI-I) was created in 2003 by Kim et al. with the purposeful of assessing the dietary esteem of diets both inside a country for observing reasons and over countries for comparative ponder. Ten components made up the DQI-I, and scores changed from to 100. Among the components were add up to balance, ampleness (natural products, vegetables, grains, fiber, protein, press, calcium, and vitamin C), and differing qualities (both generally and inside protein sources). In clinical settings, GI is utilized to assist patients select nourishments that are fitting for their blood glucose levels, particularly those who have diabetes. Boundary scores for the GI are 55 and 70. By and large talking, nourishments with a GI > 70 are considered to have a tall glycemic file. Fast-GI dinners cause a tall blood glucose top since of their rapid discharge of glucose after entering the gastrointestinal tract. Compared to high-GI dinners, suppers with a GI < 55 have a lower glycemic index and can stay within the digestive tract for a longer period of time. features a direct rate of fall, a small top esteem, a nonstop discharge of glucose, and a destitute rate of assimilation. In this manner, it is very valuable to utilize the GI to construct an perfect slim down in arrange to control and oversee human blood glucose levels.

4. Sensory Perception-Based Evaluation Techniques


In some culinary traditions, a food's "tastiness" is closely linked to its "healthiness." Although some academics contend that individuals commonly believe that solid rises to less top notch," others counter that this thought may be streamlined or that "solid = delicious" in other settings. For instance, in Chinese or French epicurean cultures, where food is prized largely for its flavor, the presumption that healthiness and tastiness are negatively correlated may not be as strongly held. If consumers don't like the way a healthy product tastes, its nutritional content is useless. Therefore, sensory factors are crucial, and a method for evaluating meals must take "taste" into account.

A novel approach to determining a food's nutritional worth may be summed up as combining dietary recommendations with sensory evaluations. New meals are created and marketed mostly using traditional sensory evaluation to satisfy customer preferences. Proper sensory analysis requires a skilled manual evaluation team. There are several methods for precisely assessing how people respond to food using the sensory test methodology. The taste wheel, for instance, is a sensory evaluation method that has been applied extensively to tea, coffee, wine, and fermented foods. For all market participants—consumers, producers, importers and exporters, merchants, tasters, and baristas—it also acts as a communication tool. The accurate sensory details of a previously assessed meal are available to consumers. Sinochem Agribusiness presented the Panda Direct, a collection of high-quality Chinese agrarian items that outlines the flavor wheel concept. It gives assessment records for rural items agreeing to a assortment of components, counting surface, taste, flavor, and appearance. This offers a methodical way to evaluate agricultural products when paired with instrument detection data. Additionally, the tasting wheel might be altered to include nutrition facts, which would help spread knowledge about diet and health.

IV. METHODS FOR FIGURING OUT HOW NUTRITIOUS A FOOD IS.


Given the extend of food-related disciplines, it is obvious that our goals may impact the most excellent assessment technique. For occasion, nourishment researchers may be more stressed approximately supplement misfortunes amid nourishment handling, indeed in spite of the fact that a Level 1 grade is satisfactory to show the dietary worth of a few items. Within the in the interim, specialists within the field of preventative medication are committed to examining the potential part that slim down and sustenance may play within the avoidance and treatment of unremitting disarranges. Hence, to affirm or explore the wellbeing impacts of nourishment, a Level 2 assessment is required. Besides, the comes

about of a dependable appraisal framework too contribute to the dispersal of nourishment information and help within the nourishment determination prepare for customers.



V. LEVELS AT WHICH FOOD'S NUTRITIONAL VALUE IS ASSESSED

The previously mentioned outline of strategies for deciding the wholesome esteem of nourishment makes it apparent that the nourishment category, evaluation objective, and inquire about profundity all have a coordinate affect on the determination of a appropriate assessment strategy. Subsequently, in arrange to urge a authentic and adequatessment, assified the sustenance assessment approaches.



Level 1: Nutrient analysis and detection are part of evaluations. they use physical and chemical analysis techniques to identify and analyse the nutrients in food, Then, compare the findings to those of a reference food or FCT to assess the nutritional value of the meal. Level 1 evaluation procedures may be used to analyse agricultural commodities, prepackaged foods, and other items since they are appropriate for most nutrition assessments and characterizing the nutritional properties of food

Level 2: assessment techniques focus on the health impacts of a product and its ingredients from a clinical standpoint, based on a Level 1 review. Various nutritional indices can indicate the biological nutritional content of food, and human

or animal experiments can further investigate and validate the health impacts. Level 2: Assessment is useful for confirming the health benefits of food and investigating its constituent parts.

Level 3: Assessments look into the mechanisms behind how particular dietary categories or nutrients impact human wellbeing, going past Level 2 assessments. In differentiate to the prior assessment strategies, a Level 3 assessment helps within the outline of components of activity and dietary physiology, as well as the examination of the physiological functions of supplements within the body at the quality and cell levels by clinical nourishment, atomic wholesome science, and other disciplines.

VI. DISCUSSION

OPPORTUNITIES TO ANALYZE FOOD NUTRITION:

Better Systems Are Needed: More advanced methods for evaluating food groups and calculating nutritional scores are required, especially when examining dietary trends. Uses in the Food Sector:

New Product Development: Targeted product development, such as low-GI meals, can be guided by nutritional assessments.

Policy Formulation: Government policies and recommendations, such as DRIs, are supported by nutritional assessments.

Consumer Guidance: Better food labeling and certification programs that are founded on precise nutritional analyses enable customers to make better decisions. Front-of-Package (FOP) systems, such as those seen in many nations, and China's "healthy choice" emblem are two examples.

Precision Food: Future food research should create "precision food" that is suited to each person's needs while taking flavor, function, and nutrients into account, taking inspiration from precision medicine.

Sustainability: Environmental effect analyses should be a part of future food evaluations, taking into account things like production-related pollution and carbon emissions. **Comprehensive Assessment:** A larger, more varied evaluation system should include nutritional value assessments.

CURRENT ISSUES AND IMPROVEMENTS:

The multidisciplinary nature of nutritional assessment Agricultural science, food science, nutrition, and even economics are all connected by research on food nutritional value. Because of its multidisciplinary character, it necessitates carefully weighing different viewpoints.

Evolution of Agricultural Focus: The nutritional quality of agricultural goods is being emphasized as Chinese agriculture moves toward green development and nutrition sensitivity.

Food business Transformation: Nutritional value and the identification of certain nutrients and advantageous components are becoming more important in the food business than flavor. Equipment modifications, packaging, and processing all play a part.

Nutrition and Disease: The necessity of balanced nutrition and dietary practices is highlighted by the ongoing research into the relationship between nutrients and chronic illnesses.

- **Limitations of Simple Nutrient Content Assessment:** Nutritional value cannot be ascertained just by measuring nutrient content. Additionally important are bioavailability (absorption and use), processing, and storage. Standardized evaluation techniques are necessary. Particularly for health foods and specialty dietary items, a uniform and rational process is required to direct the evaluation of nutritional content. The methods used nowadays differ based on the resource and objectives of the research.
- **Medical Nutrition Assessment as an Inspiration:** The article recommends using medical nutrition assessment techniques as a model to create comparable food evaluation scales and techniques. This highlights the necessity of reliable, unbiased, and valid methods specific to the food under study.

Communication of Results: In addition to following legal requirements, evaluation results should be communicated to customers in a clear and intelligible manner.

VII. CONCLUSION

- For scientific research and consumer comprehension, a thorough and uniform method of assessing the nutritional content of food is essential. Researchers from a variety of fields can benefit from the suggested three-level assessment framework, which covers a wide range of nutritional traits. This framework guarantees more precise and pertinent evaluations of food's nutritional content, suited to particular study goals, by offering a systematic process for choosing suitable assessment techniques.
- In the end, this results in a better comprehension of how food affects health and gives customers the ability to make knowledgeable dietary decisions, especially in light of the growing popularity of nutraceuticals and their possible health advantages. This systematic evaluation approach is critical for improving nutrition and advancing our understanding of food science.

REFERENCES

1. Tapsell L.C., Neale E.P., Satija A., Hu F.B. Foods, Nutrients, and Dietary Patterns: Interconnections and Implications for Dietary Guidelines. *Adv. Nutr. Int. Rev. J.* 2016;7:445–454. doi: 10.3945/an.115.011718. [DOI] [PMC free article] [PubMed] [Google Scholar]
2. Sharma I.K., Di Prima S., Essink D., Broerse J.E.W. Nutrition-Sensitive Agriculture: A Systematic Review of Impact Pathways to Nutrition Outcomes. *Adv. Nutr. Int. Rev. J.* 2020;12:251–275. doi: 10.1093/advances/nmaa103. [DOI] [PMC free article] [PubMed] [Google Scholar]
3. Ruel M.T., Quisumbing A.R., Balagamwala M. Nutrition-sensitive agriculture: What have we learned so far? *Glob. Food Secur.* 2018;17:128–153. doi: 10.1016/j.gfs.2018.01.002. [DOI] [Google Scholar]
4. Ruel M.T., Alderman H., Maternal and Child Nutrition Study Group Nutrition-sensitive interventions and programmes: How can they help to accelerate progress in improving maternal and child nutrition? *Lancet.* 2013;382:536–551. doi: 10.1016/S0140-6736(13)60843-0. [DOI] [PubMed] [Google Scholar]
5. Mailliot M., Ferguson E.L., Drewnowski A., Darmon N. Nutrient Profiling Can Help Identify Foods of Good Nutritional Quality for Their Price: A Validation Study with Linear Programming. *J. Nutr.* 2008;138:1107–1113. doi: 10.1093/jn/138.6.1107. [DOI] [PubMed] [Google Scholar]
6. Dubois C., Tharrey M., Darmon N. Identifying foods with good nutritional quality and price for the Opticourses intervention research project. *Public Health Nutr.* 2017;20:3051–3059. doi: 10.1017/S1368980017002282. [DOI] [PMC free article] [PubMed] [Google Scholar]
7. Knorr D., Froehling A., Jaeger H., Reineke K., Schlueter O., Schoessler K. Emerging technologies in food processing. In: Doyle M.P., Klaenhammer T.R., editors. *Annual Review of Food Science and Technology. Volume 2. Annual Reviews; Palo Alto, CA, USA: 2011. pp. 203–235.* [DOI] [PubMed] [Google Scholar]
8. Granato D., Barba F.J., Kovacevic D.B., Lorenzo J.M., Cruz A.G., Putnik P. Functional foods: Product development, technological trends, efficacy testing, and Safety. In: Doyle M.P., McClements D.J., editors. *Annual Review of Food Science and Technology. Volume 11. Annual Reviews; Palo Alto, CA, USA: 2020. pp. 93–118.* [DOI] [PubMed] [Google Scholar]
9. Elleuch M., Bedigian D., Roiseux O., Besbes S., Blecker C., Attia H. Dietary fibre and fibre-rich by-products of food processing: Characterisation, technological functionality and commercial applications: A review. *Food Chem.* 2011;124:411–421. doi: 10.1016/j.foodchem.2010.06.077. [DOI] [Google Scholar]
10. Sahani S., Sharma Y.C. Advancements in applications of nanotechnology in global food industry. *Food Chem.* 2020;342:128318. doi: 10.1016/j.foodchem.2020.128318. [DOI] [PubMed] [Google Scholar]
11. Ananingsih V.K., Sharma A., Zhou W. Green tea catechins during food processing and storage: A review on stability and detection. *Food Res. Int.* 2013;50:469–479. doi: 10.1016/j.foodres.2011.03.004. [DOI] [Google Scholar]
12. Mihindukulasuriya S., Lim L.-T. Nanotechnology development in food packaging: A review. *Trends Food Sci. Technol.* 2014;40:149–167. doi: 10.1016/j.tifs.2014.09.009. [DOI] [Google Scholar]
13. Knorr D., Watzke H. Food Processing at a Crossroad. *Front. Nutr.* 2019;6:85. doi: 10.3389/fnut.2019.00085. [DOI] [PMC free article] [PubMed] [Google Scholar]
14. Martinez Steele E.M., Popkin B.M., Swinburn B., Monteiro C.A. The share of ultra-processed foods and the overall nutritional quality of diets in the US: Evidence from a nationally representative cross-sectional study. *Popul. Health Metr.* 2017;15:6. doi: 10.1186/s12963-017-0119-3. [DOI] [PMC free article] [PubMed] [Google Scholar]

15. Jones D.P., Park Y., Ziegler T.R. Nutritional Metabolomics: Progress in Addressing Complexity in Diet and Health. *Annu. Rev. Nutr.* 2012;32:183–202. doi: 10.1146/annurev-nutr-072610-145159. [DOI] [PMC free article] [PubMed] [Google Scholar]
16. Reedy J., Krebs-Smith S.M., Miller P.E., Liese A.D., Kahle L.L., Park Y., Subar A.F. Higher Diet Quality Is Associated with Decreased Risk of All-Cause, Cardiovascular Disease, and Cancer Mortality among Older Adults. *J. Nutr.* 2014;144:881–889. doi: 10.3945/jn.113.189407. [DOI] [PMC free article] [PubMed] [Google Scholar]
17. Coelho-Ravagnani C.D.F., Corgosinho F.C., Sanches F.L.F.Z., Prado C.M.M., Laviano A., Mota J.F. Dietary recommendations during the COVID-19 pandemic. *Nutr. Rev.* 2020;79:382–393. doi: 10.1093/nutrit/nuaa067. [DOI] [PMC free article] [PubMed] [Google Scholar]
18. Capuano E., Pellegrini N. An integrated look at the effect of structure on nutrient bioavailability in plant foods. *J. Sci. Food Agric.* 2018;99:493–498. doi: 10.1002/jsfa.9298. [DOI] [PubMed] [Google Scholar]
19. Adams J., Tyrrell R., White M. Do television food advertisements portray advertised foods in a 'healthy' food context? *Br. J. Nutr.* 2010;105:810–815. doi: 10.1017/S0007114510004435. [DOI] [PubMed] [Google Scholar]
20. Rabassa M., Alonso-Coello P., Casino G. Nutrimedia: A novel web-based resource for the general public that evaluates the veracity of nutrition claims using the GRADE approach. *PLoS ONE.* 2020;15:e0232393. doi: 10.1371/journal.pone.0232393. [DOI] [PMC free article] [PubMed] [Google Scholar]



INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

 9940 572 462  6381 907 438  ijirset@gmail.com



www.ijirset.com

Scan to save the contact details