



International Journal of Innovative Research in Science Engineering and Technology (IJIRSET)

(A Monthly, Peer Reviewed, Refereed, Scholarly Indexed, Open Access Journal)



Impact Factor: 8.699

Volume 14, Issue 3, March 2025

Sum of Secondary Orthogonal Bimatrices in $C_{n \times n}$

Dr.R. Surendar*, M. Muthulakshmi**

Assistant Professor, Department of Mathematics, Swami Dayananda College of Arts and Science, Manjakkudi,
Tiruvarur (Dist.), Tamil Nadu, India*

P.G. Student, Department of Mathematics, Swami Dayananda College of Arts and Science, Manjakkudi, Tiruvarur
(Dist.), Tamil Nadu, India**

ABSTRACT: Let $F \in \{R, C, H\}$. Let $\mathbf{U}_{n \times n}$ be the set of secondary unitary bimatrices in $F_{n \times n}$, and let $O_{n \times n}$ be the set of secondary orthogonal bimatrices in $F_{n \times n}$. Suppose $n \geq 2$. we show that every $A_B \in F_{n \times n}$ can be written as a sum of bimatrices in $\mathbf{U}_{n \times n}$ and of bimatrices in $O_{n \times n}$. let $A_B \in F_{n \times n}$ be given that and let $k \geq 2$ be the least integer that is a least upper bound of the singular values of A_B . When $F=C$, we show that A_B can be written as a sum of k bimatrices from $\mathbf{U}_{n \times n}$.

KEYWORDS: Orthogonal matrix, unitary matrix, bimatrix, orthogonal bimatrix, unitary bimatrix, secondary orthogonal bimatrices, secondary unitary bimatrices.

AMS classification: 15A09, 15A15, 15A57.

I. INTRODUCTION

Matrices provide a very powerful tool for dealing with linear models. Bimatrices are still a powerful and an advanced tool which can handle over one linear model at a time. Bimatrices are useful when time bound comparisons are needed in the analysis of a model. Bimatrices are of several types. We denote the space of $n \times n$ complex matrices by $C_{n \times n}$. For $A \in C_{n \times n}$, A^T, A^s, A^*, A^{-1} and $\det(A)$ denote transpose, secondary transpose, conjugate transpose, inverse and determinant of A respectively. If $AA^T = A^T A = I$ then A is an orthogonal matrix, where I is an identity matrix. If $AVA^T V = VA^T V A = I$ or $AA^s = A^s A = I$ Where V is a permutation matrix with units in its secondary diagonal, A^s is an secondary orthogonal matrix. In this paper, we study secondary orthogonal bimatrices as a generalization of secondary orthogonal matrices. Some of the properties of secondary orthogonal matrices are extended to secondary orthogonal bimatrices. Some important results of secondary orthogonal matrices are generalized to secondary orthogonal bimatrices.

II. PRELIMINARIES

Definition 2.1 [7]

A bimatrix A_B is defined as the union of two rectangular array of numbers A_1 and A_2 arranged into rows and columns. It is written as $A_B = A_1 \cup A_2$ with $A_1 \neq A_2$ (except zero and unit bimatrices) where,

$$A_1 = \begin{bmatrix} a_{11}^1 & a_{12}^1 & \cdots & a_{1n}^1 \\ a_{21}^1 & a_{22}^1 & \cdots & a_{2n}^1 \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1}^1 & a_{m2}^1 & \cdots & a_{mn}^1 \end{bmatrix} \text{ and } A_2 = \begin{bmatrix} a_{11}^2 & a_{12}^2 & \cdots & a_{1n}^2 \\ a_{21}^2 & a_{22}^2 & \cdots & a_{2n}^2 \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1}^2 & a_{m2}^2 & \cdots & a_{mn}^2 \end{bmatrix}$$

' \cup ' is just for the notational convenience (symbol) only.

Definition 2.2 [7]

Let $A_B = A_1 \cup A_2$ and $C_B = C_1 \cup C_2$ be any two $m \times n$ bimatrices. The sum D_B of the bimatrices A_B and C_B is defined as

$$D_B = A_B + C_B = (A_1 \cup A_2) + (C_1 \cup C_2) = (A_1 + C_1) \cup (A_2 + C_2)$$

Where $A_1 + C_1$ and $A_2 + C_2$ are the usual addition of matrices.

Definition 2.3 [8]

If $A_B = A_1 \cup A_2$ and $C_B = C_1 \cup C_2$ be two bimatrices, then A_B and C_B are said to be equal (written as $A_B = C_B$) if and only if A_1 and C_1 are identical and A_2 and C_2 are identical. (That is, $A_1 = C_1$ and $A_2 = C_2$).

Definition 2.4 [8]

Given a bimatrix $A_B = A_1 \cup A_2$ and a scalar λ , the product of λ and A_B written as λA_B is defined to be

$$\lambda A_B = \begin{bmatrix} \lambda a_{11}^1 & \lambda a_{12}^1 & \cdots & \lambda a_{1n}^1 \\ \lambda a_{21}^1 & \lambda a_{22}^1 & \cdots & \lambda a_{2n}^1 \\ \vdots & \vdots & \ddots & \vdots \\ \lambda a_{m1}^1 & \lambda a_{m2}^1 & \cdots & \lambda a_{mn}^1 \end{bmatrix} \cup \begin{bmatrix} \lambda a_{11}^2 & \lambda a_{12}^2 & \cdots & \lambda a_{1n}^2 \\ \lambda a_{21}^2 & \lambda a_{22}^2 & \cdots & \lambda a_{2n}^2 \\ \vdots & \vdots & \ddots & \vdots \\ \lambda a_{m1}^2 & \lambda a_{m2}^2 & \cdots & \lambda a_{mn}^2 \end{bmatrix}$$

$$= (\lambda A_1 \cup \lambda A_2).$$

That is, each element of A_1 and A_2 are multiplied by λ .

Remark 2.5 [8]

If $A_B = A_1 \cup A_2$ be a bimatrix, then we call A_1 and A_2 as the component matrices of the bimatrix A_B .

Definition 2.6 [7]

If $A_B = A_1 \cup A_2$ and $C_B = C_1 \cup C_2$ are both $n \times n$ square bimatrices then, the bimatrix multiplication is defined as, $A_B \times C_B = (A_1 C_1) \cup (A_2 C_2)$.

Definition 2.7 [7]

Let $A_B^{m \times m} = A_1 \cup A_2$ be a $m \times m$ square bimatrix.

We define $I_B^{m \times m} = I_1^{m \times m} \cup I_2^{m \times m} = I_1^{m \times m} \cup I_2^{m \times m}$ to be the identity bimatrix.

Definition 2.8 [7]

Let $A_B^{m \times m} = A_1 \cup A_2$ be a square bimatrix, A_B is a symmetric bimatrix if the component matrices A_1 and A_2 are symmetric matrices. i.e, $A_1 = A_1^T$ and $A_2 = A_2^T$.

Definition 2.9 [7]

Let $A_B^{m \times m} = A_1 \cup A_2$ be a $m \times m$ square bimatrix i.e, A_1 and A_2 are $m \times m$ square matrices. A skew-symmetric bimatrix is a bimatrix A_B for which $A_B = -A_B^T$, where $-A_B^T = -A_1^T \cup -A_2^T$ i.e, the component matrices A_1 and A_2 are skew-symmetric.

Definition 2.10[4]

A bimatrix $A_B = A_1 \cup A_2$ is said to be unitary bimatrix,

if $A_B A_B^* = A_B^* A_B = I_B$ (or) $(A_1 A_1^* \cup A_2 A_2^*) = (A_1^* A_1 \cup A_2^* A_2) = I_1 \cup I_2$.

(That is, the component matrices of A_B are unitary.)

That is, $A_B^* = A_B^{-1}$ (or) $(A_1^* \cup A_2^*) = (A_1^{-1} \cup A_2^{-1})$.

Definition 2.11 [5]

A bimatrix $A_B = A_1 \cup A_2$ is said to be orthogonal bimatrix,

if $A_B A_B^T = A_B^T A_B = I_B$ (or) $(A_1 A_1^T \cup A_2 A_2^T) = (A_1^T A_1 \cup A_2^T A_2) = I_1 \cup I_2$.

(That is, the component matrices of A_B are orthogonal.)

That is, $A_B^T = A_B^{-1}$ (or) $(A_1^T \cup A_2^T) = (A_1^{-1} \cup A_2^{-1})$.

III. SECONDARY ORTHOGONAL AND SECONDARY UNITARY BIMATRICES

Definition 3.1 [6]

A bimatrix $A_B = A_1 \cup A_2$ is said to be secondary orthogonal bimatrix, if $A_B V_B A_B^T V_B = V_B A_B^T V_B A_B = I_B$ or $A_B A_B^S = A_B^S A_B = I_B$ where V_B is a permutation bimatrix with units in its secondary diagonal.

(That is, the component matrices of A_B are secondary orthogonal.)

That is, $A_B^S = A_B^{-1}$ (or) $(A_1^S \cup A_2^S) = (A_1^{-1} \cup A_2^{-1})$.

Remark 3.2

Let $A_B = A_1 \cup A_2$ be a secondary orthogonal bimatrix. If A_1 and A_2 are square and possess the same order then A_B is called square secondary orthogonal bimatrix, and if A_1 and A_2 are of different orders then A_B is called mixed square secondary orthogonal bimatrix.

Example 3.3

$$(1) A_B = \begin{bmatrix} \frac{3}{5} & \frac{4}{5} & 0 \\ -\frac{4}{5} & \frac{3}{5} & 0 \\ 0 & 0 & 1 \end{bmatrix} \cup \begin{bmatrix} \frac{2}{3} & \frac{1}{3} & \frac{2}{3} \\ \frac{1}{3} & \frac{2}{3} & -\frac{2}{3} \\ -\frac{2}{3} & \frac{2}{3} & \frac{1}{3} \end{bmatrix} \text{ is a square secondary orthogonal bimatrix.}$$

$$(2) A_B = \begin{bmatrix} \sin \theta & \cos \theta \\ \cos \theta & -\sin \theta \end{bmatrix} \cup \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & -\sin \theta \\ 0 & \sin \theta & \cos \theta \end{bmatrix} \text{ is a mixed square secondary orthogonal bimatrix.}$$

Definition 3.4[4]

Let $A_B = A_1 \cup A_2$ be an $n \times n$ complex bimatrix. (A bimatrix A_B is said to be complex if it takes entries from the complex field). A_B^S is called a secondary unitary bimatrix if $A_B V_B A_B^* V_B = V_B A_B^* V_B A_B = I_B$ (or) $\bar{A}_B^S = A_B^{-1}$.

That is, $A_1 V_1 A_1^* V_1 \cup A_2 V_2 A_2^* V_2 = V_1 A_1^* V_1 A_1 \cup V_2 A_2^* V_2 A_2 = I_1 \cup I_2$.

Example 3.5

$$A_B = A_1 \cup A_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} i & i \\ i & -i \end{bmatrix} \cup \frac{1}{2} \begin{bmatrix} 1+i & -1+i \\ 1+i & 1-i \end{bmatrix} \text{ is a secondary unitary bimatrix.}$$

In this paper, we have determined which bimatrices (if any) in $C_{n \times n}$ can be written as a sum of secondary unitary or secondary orthogonal bimatrices. We let $\mathbf{u}_{n \times n}$ and $O_{n \times n}$ are the set of secondary unitary and secondary orthogonal bimatrices in the complex field. We begin with the following observation.

Lemma 3.6

Let n be a given positive integer. Let $G \subset F_{n \times n}$ be a group under multiplication. Then $A_B \in F_{n \times n}$ can be written as a sum of bimatrices in G if and only if for every $Q_B, P_B \in G$, the bimatrix $Q_B A_B P_B$ can be written as a sum of bimatrices in G .

Notice that both $\mathbf{u}_{n \times n}$ and $O_{n \times n}$ are groups under multiplication.

Let $\alpha_1, \alpha_2 \in F$ be given. Then lemma 3.6 guarantees that for each $Q_B \in G$, we have that $\alpha_1 Q_1 \cup \alpha_2 Q_2$ can be written as a sum of bimatrices from G if and only if $\alpha_1 I_1 \cup \alpha_2 I_2$ can be written as a sum of bimatrices from G .

Lemma 3.7

Let $n \geq 2$ be a given integer. Let $G \subset F_{n \times n}$ be a group under multiplication. Suppose that G contains $K_B \equiv \text{diag}(1, -1, \dots, -1)$ and the permutation bimatrices. Then every $A_B \in F_{n \times n}$ can be written as a sum of bimatrices in G if and only if for each $\alpha_1, \alpha_2 \in F, \alpha_1 I_1 \cup \alpha_2 I_2$ can be written as a sum of bimatrices from G .

IV. SUM OF SECONDARY ORTHOGONAL BIMATRICES IN $C_{n \times n}$

The only bimatrices in the set of all secondary orthogonal bimatrices of order 1 are ± 1 . Hence, not every element of $F_{1 \times 1}$ can be written as a sum of elements in the set of all secondary orthogonal bimatrices of order 1. In fact, only the integers can be written as a sum of elements of the set of all secondary orthogonal bimatrices of order 1.

Notice that $\mathbf{u}_1 = \{e^{i\theta_1}; \theta_1 \in R\} \cup \{e^{i\theta_2}; \theta_2 \in R\}$

Set $C_{B_2} = (C_1^n \cup C_2^n) \equiv \{e^{i\theta_1} + e^{i\beta_1}; \theta_1, \beta_1 \in R\} \cup \{e^{i\theta_2} + e^{i\beta_2}; \theta_2, \beta_2 \in R\}$.

If $\theta_1, \theta_2, \beta_1, \beta_2 \in R$ are given, then $|e^{i\theta_1} + e^{i\beta_1}| \leq 2$ and $|e^{i\theta_2} + e^{i\beta_2}| \leq 2$.

Hence, $(C_1^n \cup C_2^n) \subset (A_1^n \cup A_2^n) \equiv \{Z_1 \in C; |Z_1| \leq 2\} \cup \{Z_2 \in C; |Z_2| \leq 2\}$

We show that $(A_1^n \cup A_2^n) \subset (C_1^n \cup C_2^n)$. Let $0 \leq r_1 \leq 2$ and $0 \leq r_2 \leq 2$ are given.

Set $\beta_1 = -\theta_1, \beta_2 = -\theta_2$ and Choose θ_1 and θ_2 so that, $2 \cos \theta_1 = r_1$ and $2 \cos \theta_2 = r_2$.

Then $e^{i\theta_1} + e^{-i\theta_1} = 2 \cos \theta_1 = r_1$ and $e^{i\theta_2} + e^{-i\theta_2} = 2 \cos \theta_2 = r_2$.

If $z_1 = r_1 e^{i\delta_1}$ and $z_2 = r_2 e^{i\delta_2}$, then choose $\beta_1 = -\theta_1 + 2\delta_1; \beta_2 = -\theta_2 + 2\delta_2$, and choose θ_1, θ_2 so that $2 \cos(\theta_1 - \delta_1) = r_1$ and $2 \cos(\theta_2 - \delta_2) = r_2$.

Let $k \geq 2$ be an integer.

$$\text{Set } C_{B_k} = (C_1^k \cup C_2^k) \equiv \left\{ \sum_{j=1}^k e^{i\theta_j^1}; \theta_j^1 \in R \text{ for } j = 1, \dots, k \right\} \\ \cup \left\{ \sum_{j=1}^k e^{i\theta_j^2}; \theta_j^2 \in R \text{ for } j = 1, \dots, k \right\}$$

and set $A_{B_k} = (A_1^k \cup A_2^k) \equiv \{z_1 \in C; |z_1| \leq k\} \cup \{z_2 \in C; |z_2| \leq k\}$.

We show that for each k , we have $(A_1^k \cup A_2^k) = (C_1^k \cup C_2^k)$.

First, notice that for each k , we have $(C_1^k \cup C_2^k) \subset (A_1^k \cup A_2^k)$. We now show that $(A_1^k \cup A_2^k) \subset (C_1^k \cup C_2^k)$. If $z_1 = r_1 e^{i\beta_1}; z_2 = r_2 e^{i\beta_2}$ with $r_1, r_2, \beta_1, \beta_2 \in R$ and $r_1, r_2 \geq 0$, then $e^{i\theta_1^1} + e^{i\theta_2^1} + \dots + e^{i\theta_k^1} = r_1 e^{i\beta_1}; e^{i\theta_1^2} + e^{i\theta_2^2} + \dots + e^{i\theta_k^2} = r_2 e^{i\beta_2}$ if and only if $e^{i(\theta_1^1 - \beta_1)} + \dots + e^{i(\theta_k^1 - \beta_1)} = r_1; e^{i(\theta_1^2 - \beta_2)} + \dots + e^{i(\theta_k^2 - \beta_2)} = r_2$ (1)

Hence, $(z_1 = r_1 e^{i\beta_1}; z_2 = r_2 e^{i\beta_2}) \in (C_1^k \cup C_2^k)$ if and only if $r_1, r_2 \in (C_1^k \cup C_2^k)$.

For $(\theta_1^1, \dots, \theta_k^1; \theta_1^2, \dots, \theta_k^2) \in R$, set $f_k^1(\theta_1^1, \dots, \theta_k^1) \equiv e^{i\theta_1^1} + \dots + e^{i\theta_k^1}; f_k^1(\theta_1^1, \dots, \theta_k^1) \equiv e^{i\theta_1^1} + \dots + e^{i\theta_k^1}; f_k^2(\theta_1^2, \dots, \theta_k^2) \equiv e^{i\theta_1^2} + \dots + e^{i\theta_k^2}$.

The case $k=2$ has already been shown. Let $k=3$, and suppose $0 \leq r_1 \leq 3; 0 \leq r_2 \leq 3$ Set $(\theta_3^1, \theta_3^2) = 0$ and set $\theta_1^1 = \theta^1 = -\theta_2^1$ and $\theta_1^2 = \theta^2 = -\theta_2^2$.

Then $f_3^1(\theta_1^1, \theta_2^1, \theta_3^1) = 1 + 2\cos \theta_1; f_3^2(\theta_1^2, \theta_2^2, \theta_3^2) = 1 + 2\cos \theta_2$, and (θ_1, θ_2) may be chosen so that $0 \leq r_1 \equiv 1 + 2\cos \theta_1 \leq 3; 0 \leq r_2 \equiv 1 + 2\cos \theta_2 \leq 3$.

We use mathematical induction to show the general case. The base cases $k=2$ and $k=3$ have already been shown. Assume that $k>3$ and suppose that $(C_1^k \cup C_2^k) = (A_1^k \cup A_2^k)$.

Consider $f_{k+1}^1(\theta_1^1, \dots, \theta_k^1, \theta_{k+1}^1) \equiv e^{i\theta_1^1} + \dots + e^{i\theta_k^1} + e^{i\theta_{k+1}^1}; f_{k+1}^2(\theta_1^2, \dots, \theta_k^2, \theta_{k+1}^2) \equiv e^{i\theta_1^2} + \dots + e^{i\theta_k^2} + e^{i\theta_{k+1}^2}$.

Let $z_1 = r_1 e^{i\beta_1}$ and $z_2 = r_2 e^{i\beta_2}$ are given with $0 \leq r_1 \leq k+1, 0 \leq r_2 \leq k+1$. We show that $r_1, r_2 \in (C_1^{k+1} \cup C_2^{k+1})$.

First, we show that $(A_1^{k+1} \cup A_2^{k+1}) \subseteq (C_1^{k+1} \cup C_2^{k+1})$. If k is even, choose $\theta_3^1 = \dots = \theta_{k-1}^1 = 0; \theta_3^2 = \dots = \theta_{k-1}^2 = 0$ and $\theta_4^1 = \dots = \theta_k^1 = \pi; \theta_4^2 = \dots = \theta_k^2 = \pi$. Then $f_{k+1}^1(\theta_1^1, \dots, \theta_k^1, \theta_{k+1}^1) \equiv e^{i\theta_1^1} + \dots + e^{i\theta_k^1} + e^{i\theta_{k+1}^1}; f_{k+1}^2(\theta_1^2, \dots, \theta_k^2, \theta_{k+1}^2) = e^{i\theta_1^2} + e^{i\theta_2^2}$.

If k is odd, choose $\theta_4^1 = \dots = \theta_{k-1}^1 = 0; \theta_4^2 = \dots = \theta_{k-1}^2 = 0$ and $\theta_5^1 = \dots = \theta_k^1 = \pi; \theta_5^2 = \dots = \theta_k^2 = \pi$. Then $f_{k+1}^1(\theta_1^1, \dots, \theta_k^1, \theta_{k+1}^1) \equiv e^{i\theta_1^1} + e^{i\theta_2^1} + e^{i\theta_3^1}; f_{k+1}^2(\theta_1^2, \dots, \theta_k^2, \theta_{k+1}^2) \equiv e^{i\theta_1^2} + e^{i\theta_2^2} + e^{i\theta_3^2}$.

In both cases, notice that $(A_1^{k+1} \cup A_2^{k+1}) \subseteq (C_1^{k+1} \cup C_2^{k+1})$. Hence, we may assume further that $r_1, r_2 \geq 1$; that is, we need to show that $(r_1, r_2) \in (C_1^{k+1} \cup C_2^{k+1})$ for $1 \leq r_1 \leq k+1; 1 \leq r_2 \leq k+1$.

Choose $\theta_{k+1}^1 = 0; \theta_{k+1}^2 = 0$, so that $f_{k+1}^1(\theta_1^1, \dots, \theta_k^1, \theta_{k+1}^1) = f_k^1(\theta_1^1, \dots, \theta_k^1) + 1; f_{k+1}^2(\theta_1^2, \dots, \theta_k^2, \theta_{k+1}^2) = f_k^2(\theta_1^2, \dots, \theta_k^2) + 1$. Now, by our inductive hypothesis, the equation $f_k^1(\theta_1^1, \dots, \theta_k^1) + 1 = r_1; f_k^2(\theta_1^2, \dots, \theta_k^2) + 1 = r_2$ has a solution since $0 \leq r_1 - 1 \leq k; 0 \leq r_2 - 1 \leq k$.

Lemma 4.1

Let $k \geq 2$ be a given integer. Let $(A_1^k \cup A_2^k) \equiv \{z_1 \in C; |z_1| \leq k\} \cup \{z_2 \in C; |z_2| \leq k\}$ and let $(C_1^k \cup C_2^k) \equiv \left\{ \sum_{j=1}^k e^{i\theta_j^1}; \theta_j^1 \in R \text{ for } j = 1, \dots, k \right\} \cup \left\{ \sum_{j=1}^k e^{i\theta_j^2}; \theta_j^2 \in R \text{ for } j = 1, \dots, k \right\}$. Then $(A_1^k \cup A_2^k) = (C_1^k \cup C_2^k)$.

I. The case $\mathbf{U}_{n \times n}$

Let $\alpha_1, \alpha_2 \in C$ be given. Then there exist an integer $k \geq 2$ and $\theta_1^1, \theta_2^1, \dots, \theta_k^1; \theta_1^2, \theta_2^2, \dots, \theta_k^2 \in R$ such that $\alpha_1 = f_k^1(\theta_1^1, \dots, \theta_k^1); \alpha_2 = f_k^2(\theta_1^2, \dots, \theta_k^2)$. Now, notice that

$$\begin{aligned} (\alpha_1 I_1 \cup \alpha_2 I_2) &= (f_k^1(\theta_1^1, \dots, \theta_k^1) I_1) \cup (f_k^2(\theta_1^2, \dots, \theta_k^2) I_2) \\ &= (e^{i\theta_1^1} I_1 + \dots + e^{i\theta_k^1} I_1) \cup (e^{i\theta_1^2} I_2 + \dots + e^{i\theta_k^2} I_2) \end{aligned}$$

is a sum of matrices in $\mathbf{U}_{n \times n}$.

When $n=1$, every $\alpha_1, \alpha_2 \in C$ can be written as a sum of elements of the set of all secondary unitary bimatrices of order 1. When $n \geq 2$, Lemma 3.7 guarantees that every $(A_1 \cup A_2) \in C_{n \times n}$ can be written as a sum of matrices in $\mathbf{U}_{n \times n}$.

Lemma 4.2

Let n be a given positive integer. Then every $(A_1 \cup A_2) \in C_{n \times n}$ can be written as a sum of matrices in $\mathbf{U}_{n \times n}$.

Proof

Let $(A_1 \cup A_2) \in C_{n \times n}$ be given. We look at the number of matrices that make up the sum $(A_1 \cup A_2)$.

Let $\alpha_1, \alpha_2 \in C$ be given. If $|\alpha_1|, |\alpha_2| \leq k$ for some positive integer k , then $(\alpha_1, \alpha_2) \in (A_1^k \cup A_2^k)$. Moreover, $(\alpha_1, \alpha_2) \in (A_1^m \cup A_2^m)$ for every integer $m \geq k$.

For any such m , Lemma 4.1 guarantees that there exist $\theta_1^1, \dots, \theta_m^1; \theta_1^2, \dots, \theta_m^2 \in R$ such that $\alpha_1 = e^{i\theta_1^1} + \dots + e^{i\theta_m^1}; \alpha_2 = e^{i\theta_1^2} + \dots + e^{i\theta_m^2}$.

However, if $|\alpha_1|, |\alpha_2| < k$ then $\alpha_1, \alpha_2 \notin (A_1^k \cup A_2^k)$ and α_1, α_2 cannot be written as a sum of k elements of $\mathbf{U}_1(C)$.

Write $(A_1 \cup A_2) = (U_1 \cup U_2)(\Sigma_1 \cup \Sigma_2)(V_1 \cup V_2)$ where $(U_1 \cup U_2), (V_1 \cup V_2) \in C_{n \times n}$ are secondary unitary bimatrices and

$$\Sigma_1 = \text{diag}_B(\sigma_1^1, \dots, \sigma_n^1) \text{ with } \sigma_1^1 \geq \dots \geq \sigma_n^1 \geq 0;$$

$$\Sigma_2 = \text{diag}_B(\sigma_1^2, \dots, \sigma_n^2) \text{ with } \sigma_1^2 \geq \dots \geq \sigma_n^2 \geq 0.$$

Let k be the least integer such that $\sigma_1^1, \sigma_1^2 \leq k$. Suppose that $k \geq 2$. Then for each l , we have $\sigma_l^1, \sigma_l^2 \in (A_1^k \cup A_2^k)$. Moreover, $\sigma_l^1, \sigma_l^2 \notin (A_1^{k-1} \cup A_2^{k-1})$.

Hence, $(A_1 \cup A_2)$ cannot be written as a sum of $k-1$ secondary unitary bimatrices. However, for each l , we have $\sigma_l^1 = e^{i\theta_{l1}^1} + \dots + e^{i\theta_{lk}^1}, \sigma_l^2 = e^{i\theta_{l1}^2} + \dots + e^{i\theta_{lk}^2}$, where each $(\theta_{l1}^1, \dots, \theta_{lk}^1); (\theta_{l1}^2, \dots, \theta_{lk}^2) \in R$.

For each $t = 1, \dots, k$, set

$$(U_1^t \cup U_2^t) = \text{diag}_B(e^{i\theta_{1t}^1}, \dots, e^{i\theta_{nt}^1}) \cup \text{diag}_B(e^{i\theta_{1t}^2}, \dots, e^{i\theta_{nt}^2}).$$

Then $(U_1^t \cup U_2^t) \in C_{n \times n}$ is secondary unitary bimatrix and $\sum_{t=1}^k (U_1^t \cup U_2^t) = (\Sigma_1 \cup \Sigma_2)$. Hence, $(A_1 \cup A_2)$ can be written as a sum of k secondary unitary bimatrices.

Suppose that $k=1$. If $\sigma_n^1 = 1; \sigma_n^2 = 1$, Then $(\Sigma_1 \cup \Sigma_2) = (I_1 \cup I_2)$ and $(A_1 \cup A_2)$ is secondary unitary bimatrix. If $\sigma_n^1 \neq 1; \sigma_n^2 \neq 1$, then for each l , we have $\sigma_l^1, \sigma_l^2 \in (A_1^1 \cup A_2^1)$, and $(A_1 \cup A_2)$ can be written as a sum of two secondary unitary bimatrices.

Theorem 4.3

Let $(A_1 \cup A_2) \in C_{n \times n}$ be given. Let k be the least (positive) integer so that there exist $(U_1^1 \cup U_2^1), (U_1^2 \cup U_2^2), \dots, (U_1^k \cup U_2^k) \in \mathbf{U}_{n \times n}$ satisfying $(U_1^1 \cup U_2^1) + \dots + (U_1^k \cup U_2^k) = (A_1 \cup A_2)$.

1. If $A_1 \cup A_2$ is secondary unitary bimatrix, then $k=1$.
2. If $A_1 \cup A_2$ is not secondary unitary bimatrix and $\sigma_l^1(A_1) \cup \sigma_l^2(A_2) \leq 2$ then $k=2$.
3. If $m \geq 2$ is an integer such that $m < \sigma_l^1(A_1) \cup \sigma_l^2(A_2) \leq m+1$ then $k=m+1$.

For positive integers $m \geq k$, we have $(A_1^k \cup A_2^k) \subseteq (A_1^m \cup A_2^m)$. Hence, every $(U_1 \cup U_2) \in \mathbf{U}_{n \times n}$ can be written as a sum of two or more elements of $\mathbf{U}_{n \times n}$.

It follows that every $(A_1 \cup A_2) \in C_{n \times n}$ that can be written as a sum of k elements of $\mathbf{U}_{n \times n}$ can be written as a sum of m elements of $\mathbf{U}_{n \times n}$.

II. The case $\mathbf{O}_{n \times n}$

Let $n=2$. Let $\alpha_1, \alpha_2, \beta_1, \beta_2 \in C$ be given.

$$\text{Set } (A_1(\alpha_1, \beta_1) \cup A_2(\alpha_2, \beta_2)) \equiv \begin{bmatrix} \alpha_1 & \beta_1 \\ -\beta_1 & \alpha_1 \end{bmatrix} \cup \begin{bmatrix} \alpha_2 & \beta_2 \\ -\beta_2 & \alpha_2 \end{bmatrix} \quad (2)$$

Choose β_1, β_2 such that $\alpha_1^2 + \beta_1^2 = 1; \alpha_2^2 + \beta_2^2 = 1$ and notice that $(A_1(\pm\alpha_1, \pm\beta_1) \cup A_2(\pm\alpha_2, \pm\beta_2)) \in O_{2 \times 2}$

Set $(A_1^1 \cup A_2^1) \equiv (A_1(\alpha_1, \beta_1) \cup A_2(\alpha_2, \beta_2))$ and
 $(A_1^n \cup A_2^n) \equiv (A_1(\alpha_1, -\beta_1) \cup A_2(\alpha_2, -\beta_2))$

Then $(A_1^1 \cup A_2^1) + (A_1^n \cup A_2^n) = 2[\alpha_1 I_1^n \cup \alpha_2 I_2^n]$. Lemma 3.7 guarantees that every $(A_1 \cup A_2) \in C_{2 \times 2}$ can be written as a sum of bimatrices from $O_{2 \times 2}$.

We look at the case when $n=3$. Let $\alpha_1, \alpha_2, \beta_1, \beta_2 \in F$ be given.

$$\text{Set } [B_1(\alpha_1, \beta_1) \cup B_2(\alpha_2, \beta_2)] \equiv \begin{bmatrix} 1 & 0 & 0 \\ 0 & \alpha_1 & \beta_1 \\ 0 & -\beta_1 & \alpha_1 \end{bmatrix} \cup \begin{bmatrix} 1 & 0 & 0 \\ 0 & \alpha_2 & \beta_2 \\ 0 & -\beta_2 & \alpha_2 \end{bmatrix}, \quad (3)$$

$$\text{set } [C_1(\alpha_1, \beta_1) \cup C_2(\alpha_2, \beta_2)] \equiv \begin{bmatrix} \alpha_1 & 0 & \beta_1 \\ 0 & 1 & 0 \\ -\beta_1 & 0 & \alpha_1 \end{bmatrix} \cup \begin{bmatrix} \alpha_2 & 0 & \beta_2 \\ 0 & 1 & 0 \\ -\beta_2 & 0 & \alpha_2 \end{bmatrix} \quad (4)$$

$$\text{and set } [D_1(\alpha_1, \beta_1) \cup D_2(\alpha_2, \beta_2)] \equiv \begin{bmatrix} \alpha_1 & \beta_1 & 0 \\ -\beta_1 & \alpha_1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \cup \begin{bmatrix} \alpha_2 & \beta_2 & 0 \\ -\beta_2 & \alpha_2 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad (5)$$

Choose β_1, β_2 so that $\alpha_1^2 + \beta_1^2 = 1; \alpha_2^2 + \beta_2^2 = 1$. Then $[B_1(\pm\alpha_1, \pm\beta_1) \cup B_2(\pm\alpha_2, \pm\beta_2)], [C_1(\pm\alpha_1, \pm\beta_1) \cup C_2(\pm\alpha_2, \pm\beta_2)],$ and $[D_1(\pm\alpha_1, \pm\beta_1) \cup D_2(\pm\alpha_2, \pm\beta_2)],$ are all elements of $O_{3 \times 3}$.

$$\begin{aligned} \text{Set } & (B_1^1 \cup B_2^1) \equiv [B_1(\alpha_1, \beta_1) \cup B_2(\alpha_2, \beta_2)], \\ & (B_1^n \cup B_2^n) \equiv [B_1(-\alpha_1, \beta_1) \cup B_2(-\alpha_2, \beta_2)], \\ & (C_1^1 \cup C_2^1) \equiv [C_1(\alpha_1, \beta_1) \cup C_2(\alpha_2, \beta_2)], \\ & (C_1^n \cup C_2^n) \equiv [C_1(-\alpha_1, \beta_1) \cup C_2(-\alpha_2, \beta_2)], \\ & (D_1^1 \cup D_2^1) \equiv [D_1(\alpha_1, \beta_1) \cup D_2(\alpha_2, \beta_2)], \\ \text{And } & (D_1^n \cup D_2^n) \equiv [D_1(-\alpha_1, \beta_1) \cup D_2(-\alpha_2, \beta_2)]. \end{aligned}$$

$$\begin{aligned} \text{Then, } & (B_1^1 \cup B_2^1) - (B_1^n \cup B_2^n) + (C_1^1 \cup C_2^1) - (C_1^n \cup C_2^n) \\ & + (D_1^1 \cup D_2^1) - (D_1^n \cup D_2^n) = 2[\alpha_1 I_1^m \cup \alpha_2 I_2^m] \end{aligned}$$

Lemma 3.7 now guarantees that every $(A_1 \cup A_2) \in C_{3 \times 3}$ can be written as a sum of bimatrices in $O_{3 \times 3}$.

Let $n=2m$ be a positive even integer, and let $\delta_1, \delta_2 \in C$ be given. Choose $(A_1^1 \cup A_2^1), (A_1^n \cup A_2^n) \in O_{2 \times 2}$ so that $[(A_1^1 \cup A_2^1) + (A_1^n \cup A_2^n)] = [\delta_1 I_1^n \cup \delta_2 I_2^n]$.

Set $(E_1^1 \cup E_2^1) = (A_1^1 \cup A_2^1) \oplus \dots \oplus (A_1^1 \cup A_2^1) (m \text{ copies})$

and set $(E_1^n \cup E_2^n) = (A_1^n \cup A_2^n) \oplus \dots \oplus (A_1^n \cup A_2^n) (m \text{ copies})$.

Then $(E_1^1 \cup E_2^1), (E_1^n \cup E_2^n) \in O_{2m \times 2m}$, and $[(E_1^1 + E_1^n) \cup (E_2^1 + E_2^n)] = (\delta_1 I_1^{2m} \cup \delta_2 I_2^{2m})$

Let $n = 2m + 1 \geq 3$ be an odd integer, and let $\delta_1, \delta_2 \in C$ be given.

Choose $(A_1^1 \cup A_2^1), (A_1^n \cup A_2^n) \in O_{2 \times 2}$ so that $((A_1^1 + A_1^n) \cup (A_2^1 + A_2^n)) = (\delta_1 I_1^n \cup \delta_2 I_2^n)$.

Also, choose $(B_1^1 \cup B_2^1), (B_1^n \cup B_2^n), (C_1^1 \cup C_2^1), (C_1^n \cup C_2^n), (D_1^1 \cup D_2^1), (D_1^n \cup D_2^n) \in O_{3 \times 3}$.

Such that $[(B_1^1 - B_1^n) \cup (B_2^1 - B_2^n)] + [(C_1^1 - C_1^n) \cup (C_2^1 - C_2^n)] + [(D_1^1 - D_1^n) \cup (D_2^1 - D_2^n)] = (\delta_1 I_1^m + \delta_2 I_2^m)$

Set $(E_1^1 - E_2^1) = (A_1^1 \cup A_2^1) \oplus \dots \oplus (A_1^1 \cup A_2^1) \oplus (B_1^1 \cup B_2^1)[m - 1 \text{ copies of } (A_1^1 \cup A_2^1)]$

Set $(E_1^n \cup E_2^n) = (A_1^n \cup A_2^n) \oplus \dots \oplus (A_1^n \cup A_2^n) \oplus (-B_1^n \cup -B_2^n)[m - 1 \text{ copies of } (A_1^n \cup A_2^n)],$

Set $(E_1^m \cup E_2^m) = (I_1^{2m-2} \cup I_2^{2m-2}) \oplus (C_1^1 \cup C_2^1),$

Set $(E_1^v \cup E_2^v) = -(I_1^{2m-2} \cup I_2^{2m-2}) \oplus -(C_1^n \cup C_2^n),$

Set $(E_1^p \cup E_2^p) = (I_1^{2m-2} \cup I_2^{2m-2}) \oplus (D_1^1 \cup D_2^1),$

and set $(E_1^v \cup E_2^v) = -(I_1^{2m-2} \cup I_2^{2m-2}) \oplus -(D_1^n \cup D_2^n).$

Then each $(E_1^j \cup E_2^j) \in O_{2m+1}$, and $(E_1^1 \cup E_2^1) + \dots + (E_1^v \cup E_2^v) = (\delta_1 I_1^{2m+1} \cup \delta_2 I_2^{2m+1}).$

Hence, for every $\alpha_1, \alpha_2 \in C$ and for every integer $n \geq 2$, $(\alpha_1 I_1 \cup \alpha_2 I_2)$ can be written as a sum of bimatrices from $O_{n \times n}$. Lemma 4.2 guarantees that every $(A_1 \cup A_2) \in C_{n \times n}$ can be written as a sum of bimatrices from $O_{n \times n}$.

Theorem 4.4

Let $n \geq 2$ be a given integer. Then every $(A_1 \cup A_2) \in C_{n \times n}$ can be written as a sum of bimatrices from $O_{n \times n}$.

Proof

Suppose that $(A_1 \cup A_2) = [(Q_1^1 + Q_1^n) \cup (Q_2^1 + Q_2^n)]$, where $(Q_1^1 \cup Q_2^1), (Q_1^n \cup Q_2^n) \in O_{n \times n}$.

Then one checks that $(A_1 V_1 A_1^T V_1 \cup A_2 V_2 A_2^T V_2) = (Q_1^1 A_1^S \cup Q_2^1 A_2^S)(A_1 Q_1^{1S} \cup A_2 Q_2^{1S})$, so that $(A_1 V_1 A_1^T V_1 \cup A_2 V_2 A_2^T V_2)$ and $(V_1 A_1^T V_1 A_1 \cup V_2 A_2^T V_2 A_2)$ are similar. Theorem 13 of [3] ensures that $(A_1 \cup A_2) + (Q_1 S_1 \cup Q_2 S_2)$, where $(Q_1 \cup Q_2)$ is secondary orthogonal bimatrices and $(S_1 \cup S_2)$ is secondary symmetric bimatrices (or that $(A_1 \cup A_2)$ has a $(Q_1 S_1 \cup Q_2 S_2)$ bidecomposition).

Suppose now that has a $(Q_1 S_1 \cup Q_2 S_2)$ bidecomposition. Is it true that $(A_1 \cup A_2)$ can be written as a sum of two (complex) secondary orthogonal bimatrices?

Take the case $n=1$, and notice that every $(A_1 \cup A_2) \in C_{n \times n}$ is a scalar and has a $(Q_1 S_1 \cup Q_2 S_2)$ bidecomposition.

However, only the integers can be written as a sum of secondary orthogonal bimatrices in this case.

Lemma 4.5

Let an integer $n \geq 2$ and $0 \neq \alpha_1; 0 \neq \alpha_2 \in \mathbb{C}$ be given. If $(\alpha_1 I_1 \cup \alpha_2 I_2) = (Q_1 \cup Q_2) + (W_1 \cup W_2)$ is a sum of two bimatrices from $O_{n \times n}$, then there exists a secondary skew-symmetric bimatrices $(D_1 \cup D_2) \in C_{n \times n}$ such that $(Q_1 \cup Q_2) = \left(\frac{\alpha_1}{2} I_1 \cup \frac{\alpha_2}{2} I_2\right) + (D_1 \cup D_2)$, $(W_1 \cup W_2) = \left(\frac{\alpha_1}{2} I_1 \cup \frac{\alpha_2}{2} I_2\right) - (D_1 \cup D_2)$,

and $(D_1 V_1 D_1^T V_1 \cup D_2 V_2 D_2^T V_2) = \left[\left(1 - \frac{\alpha_1^2}{4}\right) I_1 \cup \left(1 - \frac{\alpha_2^2}{4}\right) I_2\right]$.

Conversely, if there exists a secondary skew-symmetric bimatrices $(D_1 \cup D_2) \in C_{n \times n}$ such that $(D_1 V_1 D_1^T V_1 \cup D_2 V_2 D_2^T V_2) = \left[\left(1 - \frac{\alpha_1^2}{4}\right) I_1 \cup \left(1 - \frac{\alpha_2^2}{4}\right) I_2\right]$, then $(Q_1 \cup Q_2) \equiv \left(\frac{\alpha_1}{2} I_1 \cup \frac{\alpha_2}{2} I_2\right) + (D_1 \cup D_2)$ and $(W_1 \cup W_2) \equiv \left(\frac{\alpha_1}{2} I_1 \cup \frac{\alpha_2}{2} I_2\right) - (D_1 \cup D_2)$ are in $O_{n \times n}$ and $(Q_1 \cup Q_2) + (W_1 \cup W_2) = (\alpha_1 I_1 \cup \alpha_2 I_2)$.

Proof

Let an integer $n \geq 2$ and $0 \neq \alpha_1; 0 \neq \alpha_2 \in \mathbb{C}$ be given. Suppose that $(\alpha_1 I_1 \cup \alpha_2 I_2) \in C_{n \times n}$ can be written as a sum of two secondary orthogonal bimatrices, say $(\alpha_1 I_1 \cup \alpha_2 I_2) = (Q_1 \cup Q_2) + (W_1 \cup W_2)$,

Write $(Q_1 \cup Q_2) = [a_{ij}^1 \cup a_{ij}^2] = [q_1^1 \dots q_n^1] \cup [q_1^2 \dots q_n^2]$ and $(W_1 \cup W_2) = [b_{ij}^1 \cup b_{ij}^2] = [w_1^1 \dots w_n^1] \cup [w_1^2 \dots w_n^2]$. Then $b_{ij}^1 = -a_{ij}^1$ and $b_{ij}^2 = -a_{ij}^2$ for $i \neq j$.

Now, for each $i = 1, \dots, n$, we have $\sum_{j=1}^n a_{ij}^{12} = q_i^{1T} q_i^1 = 1 = w_i^{1T} v_i^1 = \sum_{j=1}^n b_{ij}^{12} = b_{ii}^{12} + \sum_{j \neq i, j=1}^n a_{ij}^{12}$

and $\sum_{j=1}^n a_{ij}^{22} = q_i^{2T} q_i^2 = 1 = w_i^{2T} v_i^2 = \sum_{j=1}^n b_{ij}^{22} = b_{ii}^{22} + \sum_{j \neq i, j=1}^n a_{ij}^{22}$

Hence, $b_{ii}^1 = \pm a_{ii}^1$ and $b_{ii}^2 = \pm a_{ii}^2$.

Because $(Q_1 \cup Q_2) + (W_1 \cup W_2) = (\alpha_1 I_1 \cup \alpha_2 I_2)$ and $\alpha_1 \neq 0; \alpha_2 \neq 0$ we have $b_{ii}^1 = a_{ii}^1 = \frac{\alpha_1}{2}; b_{ii}^2 = a_{ii}^2 = \frac{\alpha_2}{2}$. Set $(D_1 \cup D_2) = [d_{ij}^1 \cup d_{ij}^2]$, with $d_{ij}^1 = a_{ij}^1; d_{ij}^2 = a_{ij}^2$ if $i \neq j$, and $d_{ii}^1 = 0; d_{ii}^2 = 0$, so that $(Q_1 \cup Q_2) = \left(\frac{\alpha_1}{2} I_1 \cup \frac{\alpha_2}{2} I_2\right) + (D_1 \cup D_2)$ and $(W_1 \cup W_2) = \left(\frac{\alpha_1}{2} I_1 \cup \frac{\alpha_2}{2} I_2\right) - (D_1 \cup D_2)$.

Now, since $(Q_1 \cup Q_2)$ and $(V_1 \cup V_2)$ are secondary orthogonal bimatrices, we have

$$(Q_1 V_1 Q_1^T V_1 \cup Q_2 V_2 Q_2^T V_2) \equiv \left(\frac{\alpha_1^2}{4} I_1 \cup \frac{\alpha_2^2}{4} I_2\right) + \left[\frac{\alpha_1}{2} (D_1 + D_1^T) \cup \frac{\alpha_2}{2} (D_2 + D_2^T)\right] + (D_1 V_1 D_1^T V_1 \cup D_2 V_2 D_2^T V_2) \\ = (I_1 \cup I_2) \quad (6)$$

$$\text{And } (W_1 W_1^S \cup W_2 W_2^S) \equiv \left(\frac{\alpha_1^2}{4} I_1 \cup \frac{\alpha_2^2}{4} I_2\right) - \left[\frac{\alpha_1}{2} (D_1 + D_1^T) \cup \frac{\alpha_2}{2} (D_2 + D_2^T)\right] + (D_1 V_1 D_1^T V_1 \cup D_2 V_2 D_2^T V_2) \\ = (I_1 \cup I_2) \quad (7)$$

Subtracting equation (7) from equation (6), we get $(D_1 \cup D_2) = -(D_1^S \cup D_2^S)$, so that $(D_1 \cup D_2)$ is secondary skew-symmetric bimatrices.

Moreover $(D_1 V_1 D_1^T V_1 \cup D_2 V_2 D_2^T V_2) = \left[\left(1 - \frac{\alpha_1^2}{4}\right) I_1 \cup \left(1 - \frac{\alpha_2^2}{4}\right) I_2\right]$.

For the converse, suppose that $(D_1 \cup D_2) \in C_{n \times n}$ is secondary skew-symmetric bimatrices and satisfies, $(D_1 V_1 D_1^T V_1 \cup D_2 V_2 D_2^T V_2) = \left[\left(1 - \frac{\alpha_1^2}{4}\right) I_1 \cup \left(1 - \frac{\alpha_2^2}{4}\right) I_2\right]$.

Set $(Q_1 \cup Q_2) \equiv \left(\frac{\alpha_1}{2} I_1 \cup \frac{\alpha_2}{2} I_2\right) + (D_1 \cup D_2)$ and set $(W_1 \cup W_2) \equiv \left(\frac{\alpha_1}{2} I_1 \cup \frac{\alpha_2}{2} I_2\right) - (D_1 \cup D_2)$.

Then one checks that both $(Q_1 \cup Q_2)$ and $(W_1 \cup W_2)$ are secondary orthogonal bimatrices and $[(Q_1 + W_1) \cup (Q_2 + W_2)] = (\alpha_1 I_1 \cup \alpha_2 I_2)$.

Remark 4.6

When $\alpha_1 = 0; \alpha_2 = 0$, then for any secondary orthogonal bimatrix $(Q_1 \cup Q_2)$, notice that $(\alpha_1 I_1 \cup \alpha_2 I_2) = [(Q_1 + (-Q_1)) \cup (Q_2 + (-Q_2))]$ is a sum of two secondary orthogonal bimatrices.

Let $n=2$ and $\alpha_1 \neq 0; \alpha_2 \neq 0$. Set $\beta_1 \equiv \sqrt{1 - \frac{\alpha_1^2}{4}}; \beta_2 \equiv \sqrt{1 - \frac{\alpha_2^2}{4}}$ (either square root).

Then $(D_1 \cup D_2) \equiv \begin{bmatrix} 0 & \beta_1 \\ -\beta_1 & 0 \end{bmatrix} \cup \begin{bmatrix} 0 & \beta_2 \\ -\beta_2 & 0 \end{bmatrix}$ is secondary skew-symmetric bimatrix and satisfies

$$(D_1 V_1 D_1^T V_1 \cup D_2 V_2 D_2^T V_2) = \left[\left(1 - \frac{\alpha_1^2}{4}\right) I_1 \cup \left(1 - \frac{\alpha_2^2}{4}\right) I_2 \right].$$

Lemma 4.5 guarantees that $(\alpha_1 I_1 \cup \alpha_2 I_2)$ can be written as a sum of two secondary orthogonal bimatrices.

If $n=2k$ and $\alpha_1 \neq 0; \alpha_2 \neq 0$, set $(E_1 \cup E_2) = (D_1 \cup D_2) \oplus \dots \oplus (D_1 \cup D_2)$ (k copies) and notice that $(E_1 \cup E_2)$ is secondary skew-symmetric bimatrix and satisfies $(E_1 V_1 E_1^T V_1 \cup E_2 V_2 E_2^T V_2)(E_1 E_1^T \cup E_2 E_2^T) = \left[\left(1 - \frac{\alpha_1^2}{4}\right) I_1 \cup \left(1 - \frac{\alpha_2^2}{4}\right) I_2 \right]$.

Hence, if $n=2k$ and if α_1, α_2 be a scalar, then $(\alpha_1 I_1 \cup \alpha_2 I_2)$ can be written as a sum of two secondary orthogonal bimatrices.

Theorem 4.7

Let n be a given positive integer. For each $\alpha_1, \alpha_2 \in \mathbb{C}$ and each secondary orthogonal bimatrix $(Q_1 \cup Q_2) \in C_{2n}$, $(\alpha_1 Q_1 \cup \alpha_2 Q_2)$ can be written as a sum of two secondary orthogonal bimatrices.

Remark 4.8

Let an integer $n \geq 2$ be given. If $\alpha_1, \alpha_2 \in \{-2, 0, 2\}$ then one checks that $(\alpha_1 I_1 \cup \alpha_2 I_2) \in C_{n \times n}$ can be written as a sum of two secondary orthogonal bimatrices.

Theorem 4.9

Let $\alpha_1, \alpha_2 \in \mathbb{C}$ and let a positive integer n be given. Then $(\alpha_1 I_1 \cup \alpha_2 I_2) \in C_{2n+1}$ can be written as a sum of two secondary orthogonal bimatrices from $O_{n \times n}$ if and only if $\alpha_1, \alpha_2 \in \{-2, 0, 2\}$.

Proof

For the forward implication, let $\alpha_1, \alpha_2 \in \mathbb{C}$ and let a positive integer n be given.

Suppose that $(\alpha_1 I_1 \cup \alpha_2 I_2) \in C_{2n+1}$ can be written as a sum of two secondary orthogonal bimatrices.

Then $\alpha_1 = 0; \alpha_2 \neq 0$ (or) $\alpha_1 \neq 0; \alpha_2 = 0$. If $\alpha_1 = 0; \alpha_2 = 0$, then $\alpha_1, \alpha_2 \in \{-2, 0, 2\}$.

If $\alpha_1 \neq 0; \alpha_2 \neq 0$, we show that $\alpha_1 = \alpha_2 = \pm 2$. Lemma 3.5 guarantees that there exists a secondary skew-symmetric bimatrix $(D_1 \cup D_2) \in C_{n \times n}$ satisfying

$$(D_1 V_1 D_1^T V_1 \cup D_2 V_2 D_2^T V_2) = \left[\left(1 - \frac{\alpha_1^2}{4}\right) I_1 \cup \left(1 - \frac{\alpha_2^2}{4}\right) I_2 \right].$$

Now, since n is odd, the secondary skew-symmetric bimatrix $(D_1 \cup D_2)$ is singular. Hence, $(D_1 V_1 D_1^T V_1 \cup D_2 V_2 D_2^T V_2)$ is singular and $\alpha_1 = \alpha_2 = \pm 2$.

The backward implication can be show by direct computation.

V. CONCLUSION

Here, we have studied some properties and important results of secondary orthogonal bimatrices as a generalization of orthogonal and secondary orthogonal matrices. Also, we have determined which bimatrices (if any) in $C_{n \times n}$ can be written as a sum of secondary unitary or secondary orthogonal bimatrices.

REFERENCES

- [1] Horn, R.A. and Johnson, C.R., "Matrix Analysis", Cambridge University Press, New York, 1985.
- [2] Horn, R.A. and Johnson, C.R., "Topics in Matrix Analysis", Cambridge University Press, New York, 1991.
- [3] Horn, R.A. and Merino, D.I., "Contragredient equivalence: a canonical form and some applications", *Linear Algebra Appl.*, 214 (1995), 43-92.

- [4] Ramesh, G. and Maduranthaki, P., “On Unitary Bimatrices”, *International Journal of Current Research*, Vol. **6**, Issue 09, September 2014, pp. 8395-8407.
- [5] Ramesh, G., Jothivasan, S., Muthugobal, B.K.N. and Surendar, R., “On orthogonal bimatrices”, *International Journal of Applied Research*, 1(11); 2015, pp. 1013-1024.
- [6] Surendar, R. and Renuga Saraswathi., “On secondary orthogonal bimatrices”, *International Journal of Innovative Research in Science, Engineering and Technology*, Vol. **13**, Issue 2, February 2024, pp. 909-922.
- [7] Vasantha Kandasamy, W. B., Florentin Samarandache and Ilanthendral, K., “Introduction to Bimatrices.” 2005.
- [8] Vasantha Kandasamy, W. B., Florentin Samarandache and Ilanthendral, K., “Applications of bimatrices to some Fuzzy and Neutrosophic models.” *Hexis, Phonix, Arizone*, 2005.



INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

 9940 572 462  6381 907 438  ijirset@gmail.com



www.ijirset.com

Scan to save the contact details