




International Journal of Innovative Research in Science Engineering and Technology (IJIRSET)

(A Monthly, Peer Reviewed, Refereed, Scholarly Indexed, Open Access Journal)



Impact Factor: 8.699

Volume 14, Issue 3, March 2025

The Impact of Climatic Changes on Cotton Crop Productivity: Present Scenario

Priyanka, Dr. Neetu Singh

Research Scholar, Department of Zoology, Shri Jagdishprasad Jhabarmal Tibrewala University, Jhunjhunu,
Rajasthan, India

Assistant Professor, Department of Zoology, Shri Jagdishprasad Jhabarmal Tibrewala University, Jhunjhunu,
Rajasthan, India

ABSTRACT: Climate change has a profound impact on cotton crop productivity, presenting significant challenges to global cotton production. This study explores the effects of climatic variations, such as temperature fluctuations, irregular rainfall, and the growing occurrence of extreme weather events, on cotton growth and yield. Higher temperatures can reduce the duration of the cotton growing season, while inconsistent precipitation patterns contribute to either water stress or excessive moisture, both of which adversely affect crop health. Moreover, rising CO₂ levels influence pest behavior and soil nutrient dynamics, further shaping cotton productivity. Gaining insights into these changes is essential for developing effective adaptation strategies to promote sustainable cotton cultivation amid ongoing climate shifts.

KEYWORDS: Climate change, Global Warming, Rainfall Patterns, Cotton Crop Productivity, Heat Stress.

I. INTRODUCTION

Cotton is an essential cash crop cultivated worldwide for its natural fiber, which is a key raw material in the textile industry. Originally grown in tropical and subtropical regions, its cultivation has expanded to major cotton-producing countries such as the United States, India, China, and Brazil. As a member of the Malvaceae family, cotton thrives in warm climates with extended frost-free periods, abundant sunlight, and moderate rainfall, making these conditions ideal for its growth and fiber development.

Cotton is predominantly cultivated in well-drained, loamy soils, requiring specific conditions for optimal growth. The farming process involves multiple stages, including land preparation, sowing, irrigation, pest control, and harvesting. To enhance resistance against pests and diseases, many modern cotton varieties have been genetically modified, leading to improved yields. However, cotton cultivation is highly resource-intensive, particularly in terms of water consumption and pesticide application, raising environmental concerns. To address these challenges, sustainable farming methods such as organic cultivation and advanced water management techniques are being increasingly adopted. As a crucial crop in the global economy, cotton serves as a primary raw material for the textile industry and provides employment opportunities for millions worldwide.

II. REVIEW OF LITERATURE

Aman Deep Ranga and Vasudha kak (2020) Cotton crop outputs are the major natural fiber in the world and as well as oilseed crops (Constable and Bange, 2015). While cotton is mostly grown in 80 countries, India leads in cotton production followed by China, United States, Pakistan and Brazil (Khan et al., 2020). Malvaceae family includes Gossypium species which comprises of 90 genera, including the genus Gossypium. Species of this genus shows development of bushes and saplings up to seven feet tall with an inclination of whole leaves rotating with International Journal of Current Microbiology and Applied Sciences.

D. S. Jadhav, et al., (2020) The production and Productivity of cotton crop in India was Positive and statistically significant (i.e., 2.20, 4.84 and 2.78 per cent respectively). Coefficient of variation for area was very Low as compared

to production and yield. During the study period, instability in cotton Production is higher than the instability in the area and similar to yield at all India Level.

J.B. Naveena and Sharana Basappa (2019) Cotton is the most important cash crop of monsoon season in India and popularly known as “white gold”. India has a unique distinction of being the only country in the world to cultivate all four cultivable *Gossypium* species. *Gossypium hirsutum* represents 99.9 per cent of the hybrid cotton in India and all the current Bt cotton hybrids are either *G. hirsutum* or inter-specific hybrids with *G. barbadense*. In India, cotton is grown under 40 per cent irrigated and 60 per cent rainfed. American cotton is highly susceptible to sucking insect pests and of which jassid, *A. biguttula* also referred, as leafhopper is important sucking pest. Although introduction of Bt cotton could reduce the bollworm incidence, number of other pests viz., leafhopper, mirid bugs, aphids and thrips are assuming potential threats. At present many sucking pests of cotton have assumed potential threats and causing significant economic yield losses in cotton. The reduction of pyrethroid and other conventional insecticides on Bt cotton is presumed to have enhanced infestation of sucking pests.

S. K. Muhammad, et al., (2014) Cotton aphid has a major impact on quality and yield of cotton which emphasizes the need to manage this notorious pest. The main goal of this review is to highlight various strategies viz., biological, chemical and cultural control for cotton aphid management. The selection of suitable control strategy is made on by viewing the severity of cotton aphid outbreak. The preservation of the cotton aphid natural enemies could be an ecologically sustainable method of maintaining the aphid population below to the threshold level.

III. CLIMATE CHANGE AND ITS KEY FACTORS AFFECTING COTTON PRODUCTIVITY

Climate change has significantly altered the environmental conditions required for optimal cotton growth, affecting various physiological and agronomic factors. Several key climate variables influence cotton productivity, including temperature variations, rainfall patterns, soil moisture availability, CO₂ levels, and pest and disease dynamics.

3.1. Temperature Variations

Temperature plays a critical role in different growth stages of the cotton plant, including germination, vegetative growth, flowering, and boll development. Rising temperatures can accelerate crop maturation, potentially reducing the duration of fiber development and affecting fiber quality. Extreme heat stress, particularly during the flowering and boll formation stages, leads to flower shedding, reduced boll retention, and lower yields. Additionally, heat stress increases the rate of respiration, resulting in greater energy loss and lower carbohydrate accumulation in cotton plants. Prolonged exposure to high temperatures can also affect enzyme activity, leading to metabolic imbalances and reduced crop resilience.

3.2. Rainfall Patterns

Unpredictable rainfall patterns, including excessive precipitation and prolonged drought conditions, pose significant challenges to cotton cultivation. Excessive rainfall can lead to waterlogging, which suffocates roots, reduces nutrient uptake, and promotes fungal diseases such as root rot and boll rot. Conversely, drought conditions limit soil moisture availability, restricting plant growth, causing premature leaf senescence, and reducing boll size and fiber quality. Inconsistencies in precipitation also affect the timing of irrigation schedules, increasing the dependence on artificial water sources and raising production costs.

3.3. Soil Moisture and Water Availability

Cotton is highly sensitive to soil moisture fluctuations, and its productivity is directly linked to adequate and timely water supply. Climate change-induced shifts in precipitation patterns impact soil moisture retention and irrigation requirements. Water scarcity due to prolonged dry spells reduces the plant's ability to absorb essential nutrients, weakening growth and yield. Additionally, erratic rainfall may result in soil erosion, loss of organic matter, and declining soil fertility, further affecting cotton production. Efficient water management strategies, such as drip irrigation and rainwater harvesting, are necessary to ensure sustainable water use in cotton farming.

3.4. CO₂ Levels and Photosynthesis

Rising atmospheric CO₂ concentrations can influence photosynthesis, plant metabolism, and fiber development in cotton. Studies suggest that elevated CO₂ levels may enhance photosynthetic efficiency, leading to increased biomass

production and improved water-use efficiency. However, the benefits of higher CO₂ levels may be offset by other climate-related stressors, such as heat and drought. The interaction between CO₂ enrichment and temperature variations needs further research to determine its overall impact on cotton fiber yield and quality.

3.5. Pest and Disease Incidences

Changes in temperature and humidity have altered pest and disease dynamics in cotton fields, increasing the prevalence of insect pests such as bollworms, aphids, and whiteflies. Warmer conditions accelerate pest life cycles, leading to higher infestation rates and increased pesticide dependency. Additionally, climate change favors the spread of fungal, bacterial, and viral diseases that weaken cotton plants and reduce yields. The emergence of pesticide-resistant pests due to excessive chemical use further complicates pest management, necessitating the adoption of integrated pest management (IPM) strategies.

IV. RESEARCH METHODOLOGY

The methodology outlined will enable a comprehensive evaluation of the effects of climate change on cotton crop productivity, providing insights that are globally relevant yet sensitive to local contexts. The findings will contribute to informed decision-making in agricultural policy and climate adaptation strategies.

1. Data Collection Methods

- **Primary Data:**
 - **Surveys and Questionnaires:**
 - **Target Participants:** Cotton farmers, agricultural experts, and climate scientists.
 - **Content:** Questions related to changes in temperature, rainfall patterns, and extreme weather events; the impact of these changes on cotton yield, quality, and farming practices.
 - **Field Experiments:**
 - Conduct experiments in cotton fields to observe the effects of controlled climatic variables (e.g., temperature, water availability) on crop growth and yield.
- **Secondary Data:**
 - **Historical Climate Data:**
 - Obtain data on temperature, precipitation, and other climatic factors from meteorological departments or climate databases.
 - **Agricultural Yield Data:**
 - Gather historical and current data on cotton yield from agricultural departments, research institutes, or databases like FAO.
 - **Remote Sensing and GIS:**
 - Use satellite imagery and GIS tools to analyze changes in land use, crop patterns, and environmental stress on cotton fields.

2. Sampling Strategy

- **Sampling Method:** Stratified random sampling to ensure representation from different cotton-growing regions.
- **Sample Size:** Determine based on the population of cotton farmers and the variability in climatic conditions across different regions.

3. Data Analysis

- **Quantitative Analysis:**
 - **Statistical Analysis:**
 - Use statistical tools (e.g., regression analysis, ANOVA) to assess the relationship between climatic variables and cotton productivity.
 - **Trend Analysis:**
 - Analyse trends over time to understand the long-term impact of climate change on cotton yield.
- **Qualitative Analysis:**
 - **Content Analysis:**

- Analyse responses from surveys and interviews to identify common themes related to the impact of climate change on cotton farming practices.
- Geospatial Analysis:
 - Use GIS software to map and visualize the spatial impact of climate change on cotton-growing regions.
- 4. Validation and Reliability
 - Validation Methods:
 - Cross-check data with multiple sources.
 - Use triangulation to validate findings by comparing results from different data sources and methods.
 - Reliability Testing:
 - Perform reliability tests (e.g., Cronbach's alpha) on survey instruments to ensure consistency.

V. LIMITATIONS

- Geographical Limitations: The study may be limited to specific regions due to accessibility or resource constraints.
- Data Limitations: Availability and accuracy of historical climate and yield data might affect the analysis.
- Temporal Limitations: The study may be constrained by the time frame in which data is collected, especially in the context of long-term climate trends.

VI. SUMMARY

Climate change presents substantial challenges to global agricultural productivity, including the cultivation of cotton, a key cash crop. This study explores the impact of climate change on cotton production across major cotton-growing regions such as the United States, India, China, and Brazil. Rising temperatures, shifting precipitation patterns, and the increasing occurrence of extreme weather events are significantly influencing cotton yields. Elevated temperatures can shorten the crop's growth cycle, potentially affecting fiber quality and reducing overall yield. Variability in rainfall can result in either drought stress or waterlogging, both of which adversely affect cotton productivity.

By utilizing climate data, crop simulation models, and geospatial analysis, this research identifies existing trends and forecasts potential future scenarios for cotton production under changing climate conditions. Additionally, insights from farmers and agricultural experts highlight adaptive strategies being implemented, including adjustments in planting schedules, the use of drought-resistant cotton varieties, and enhanced irrigation techniques. These findings underscore the necessity for climate-resilient agricultural practices to ensure sustainable cotton production in the face of ongoing environmental changes.

The impact of climate change on cotton productivity is evident through multiple environmental stressors, including temperature fluctuations, erratic rainfall, water scarcity, altered CO₂ levels, and increasing pest and disease pressure. Addressing these challenges requires the adoption of climate-resilient agricultural practices, improved irrigation techniques, breeding for heat- and drought-tolerant cotton varieties, and sustainable pest control methods. By implementing adaptive strategies, cotton farmers can mitigate climate-related risks and ensure stable cotton production in the face of changing environmental conditions.

REFERENCES

1. **Aman, Deep, Ranga., Vasudha, Kak., Mayur, S., Darvhankar. (2020).** Genetic and Molecular Research of Resistance to Wilt in Cotton: A Concise Review. *International Journal of Current Microbiology and Applied Sciences*, 9(6):2410-2422. doi:10.20546/IJCMAS.2020.906.296
2. **Bange, M. P., & Milroy, S. P. (2004).** Impact of short-term exposure to hot temperatures on cotton flowering and fruit retention. *Field Crops Research*, 89(2-3), 187-199.
3. **DNFI (2019).** Natural Fibres and the World Economy 2019. Available at: http://www.dnfi.org/coir/natural-fibres-and-the-world-economy-july-2019_18043/Cotton made in Africa aid by Trade Foundation, 2015. Annual report 2015.

4. **Hatfield, J. L., & Prueger, J. H. (2015).** Temperature extremes: Effect on plant growth and development. *Weather and Climate Extremes*, 10, 4-10.
5. **J.B., Naveena., Sharana, basappa. (2019).** Screening of Different Bt and Non Bt Cotton Hybrids against Aphids Aphis gossypii, and Whiteflies Bemisia tabaci on Bt Cotton. *International Journal of Current Microbiology and Applied Sciences*, 8(10): 2365-2368.doi: 10.20546/IJCMAS.2019.810.273
6. **IPCC (Intergovernmental Panel on Climate Change). (2021).** Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the IPCC. *Cambridge University*
7. **Lobell, D. B., Schlenker, W., & Costa-Roberts, J. (2011).** Climate trends and global crop production since 1980. *Science*, 333(6042), 616-620.
8. **Singh, R. P., Singh, D., & Reddy, K. R. (2007).** Effects of high temperature and elevated CO₂ on cotton growth and productivity. *Plant Science*, 173(6), 784-795. <https://doi.org/10.1016/j.plantsci.2007.09.002>
9. **Zhao, G., Bryan, B. A., Song, X., & King, D. (2015).** Impacts of climate change on cotton production in China and adaptation strategies. *Climatic Change*, 132(4), 545-558.



INTERNATIONAL
STANDARD
SERIAL
NUMBER
INDIA



INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH

IN SCIENCE | ENGINEERING | TECHNOLOGY

 9940 572 462  6381 907 438  ijirset@gmail.com



www.ijirset.com

Scan to save the contact details